

Figure 3.3 Influence of tidal range on seagrass depth distribution. Minimum depth is deeper with greater tidal range (Koch 2001). Reprinted with permission from Estuarine Research Federation.

tidal amplitude (Figure 3.3) and wave energy (Figure 3.4) as two factors that influence seagrass depth distribution. Tolerances to common abiotic factors may differ between populations of a species from different environments, as shown for R. maritima grown from seed (Koch and Dawes 1991b). Different strains grown from seeds collected in the Weeki Wachee River estuary and from North Carolina had growth rates that were similar under different combinations of temperature and photoperiod but differed with regard to salinity. This was also reported in earlier studies for other species (McMillan 1979, 1984). In those studies, different strains of T. testudinum, S. filiforme, and H. wrightii showed different tolerances to temperature, and these tolerances were often related to distinct environmental origins. These findings suggest that transplanting from one area to a different site at greater distances (e.g., seeds from Biscayne Bay to Tampa Bay) may not be as successful as when using local plants.

A large global-level literature base shows the critical effects of water quality on the presence, growth, and vitality of seagrasses. Adequate light transmission is vital to the presence of seagrasses and affects the density and biomass of seagrasses at all depths (Dixon and Leverone 1991; Duarte 1991; Johansson 1991; Kenworthy and Haunert 1991; Lapointe and Clark 1992; Tomasko 1992, 2002; Lapointe et al. 1994; Fletcher and Fletcher 1995; Kenworthy and Fonseca 1996; Kurz et al. 1999, 2000; Dixon 2000; Neely 2000). Depth determinations (±10 cm) can be obtained with a mapping-grade differential Global Positioning System, carrier-phase equipment and surveyed benchmarks (Johansson 2002b). In Tampa Bay, the deep edges for H. wrightii ranged from -0.30 to -0.76 m below mean low water (MLW), from -1.19 to




Figure 3.4 Influence of wave energy on seagrass depth distribution. Greater wave energy and deeper wave-mixing depths result in deeper minimum depths (Koch 2001). Reprinted with permission from Estuarine Research Federation.

−1.96 m for *S. filiforme*, and from −0.57 to −1.73 m for *T. testudinum*. The maximum depth of *T. testudinum* at Port Manatee in Tampa Bay was −1.6 m, in agreement with the above determinations carried out before the 1997–1998 *El Niño* event (Dixon and Leverone 1995). The shallowest deep edges for all three species occurred at the northernmost sites, and the deepest deep edges were near the mouth of Tampa Bay, where more oceanic (clear water) conditions exist.

Neely (2000) demonstrated an inverse relationship between light reduction and short-shoot numbers for H. wrightii in Tampa Bay. Leaves on plants exposed to a 43% light reduction were longer than those on shoots in control plots or in other lightreduction treatments. Her experiments showed that loss of biomass was a result of winter die-back and also occurred with a >60% reduction in light. The data showed that epiphytes and surface scattering of light must be accounted for when calculating carbon budgets. Durako and Hall (1992) found that for T. testudinum the carbon supply became nonlimiting when light was reduced to levels that limited photosynthetic rates. Tomasko (1992) demonstrated that H. wrightii, when growing under a canopy of T. testudinum leaves, had longer rhizome internodes, less frequent branching, and lower above-ground to below-ground biomass ratios than when growing alone. Studies by Sand-Jensen (1977, 1990), Cambridge et al. (1986), and Tomasko and Lapointe (1991) showed that heavy epiphyte loads on seagrass leaves are indicative of nutrient over-loading in the water column and, therefore, eutrophication, i.e., poor water quality (see Chapter 5). In addition to the adverse effects of nutrient-enriched water on photosynthetic functioning of seagrass leaves, increased levels of nutrients can result in high epiphyte loads that shade and weigh down the leaves, further lowering them in the water column.

Table 3.1 Light requirements of Florida seagrasses.

	Minimum Light Requirement	
Species and Sites	(percentage of total surface irradiance)	Reference
Thalassia testudinum		
Big Bend area, FL	15.3%	Iverson and Bittaker (1986), as reported by Duarte (1991) and by Dennison <i>et al.</i> (1993)
Northwest Cuba	23.5%	Duarte (1991)
Puerto Rico	24.4%	Duarte (1991)
Charlotte Harbor, FL	20%	Tomasko and Hall (1999)
Tampa Bay, FL	22.5%	Dixon and Leverone (1995)
Tampa Bay, FL	20.1% (annual mean)	Dixon (2000)
Corpus Christi Bay, TX	14% (4% survival after 12 months)	Czerny and Dunton (1995)
Syringodium filiforme		
Florida Bay, FL	18.3%	Iverson and Bittaker (1986) as reported by Duarte (1991)
Indian River Lagoon, FL	17.2%	Duarte (1991) pers. comm. from J. Kenworthy (1990)
Indian River Lagoon, FL (southern region)	24%–37%	Kenworthy and Fonseca (1996)
Halodule wrightii		
Indian River Lagoon, FL	17.2%	Dennison <i>et al.</i> (1993) <i>pers. comm.</i> from J. Kenworthy (1990)
Florida Bay, FL	10%-20%	Fourqurean and Zieman (1991)
Laguna Madre, TX	7%–23%	Tomasko and Dunton (1991); Dunton (1994); Czerny and Dunton (1995)

The relationship between light, both quantity and quality, and the depth at which seagrasses grow was the subject of two workshops in Florida in the 1990s, one sponsored by NOAA (Kenworthy and Haunert 1991) and the other by the Indian River Lagoon National Estuary Program (Morris and Tomasko 1993). Even though they can grow under high irradiences, seagrasses are regarded as shade plants (Hillman et al. 1989), with compensation values (I_c) of 25 to 50 µmole-photons m⁻² s⁻¹ and saturation irradiances (I_k) of <200 µmolephotons m⁻² s⁻¹. Studies comparing photosynthetic responses of seagrasses to different irradiance levels (i.e., Ic, Ik, and Pmax levels) often show a relationship to temperature and salinity, as seen for two species of Halophila (Dawes et al. 1989). For example, the estuarine, intertidal populations of H. johnsonii in the Indian River Lagoon tolerate a wide range of temperatures (10°, 20°, and 30°C) and salinities (15, 25, and 35 ppt) and did not become photoinhibited by high photosynthetically active radiation (PAR; ca. 500 µmole-photons m⁻² s⁻¹). In contrast, *H. decipiens* from deeper, nonestuarine Gulf-coast populations were intolerant of a 10°C temperature, 15 and 25 ppt salinity, and PAR above 300 µmole-photons m⁻² s⁻¹ (Dawes *et al.* 1989).

Depth distributions of Florida's seagrasses are governed by PAR levels in Tampa Bay (Miller and McPherson 1995; Dixon and Leverone 1995, 1997; Dixon 2002) and the Indian River Lagoon (Gallegos and Kenworthy 1996). Recent work has demonstrated more specific light requirements for the dominant seagrass species (Table 3.1). PAR levels measured at the deep edges of T. testudinum beds in Tampa Bay were significantly higher than the required average light level of 11% (of surface value) proposed for seagrasses by Duarte (1991). Also, T. testudinum in Florida Bay (Fourqurean and Zieman 1991) and Caribbean seagrasses (Onuf 1994) required at least 22% of surface PAR levels, when whole-plant requirements and average daily irradiances were considered. Continuous in situ measurements of underwater PAR levels for one year at the shallow and deep edges of four T. testudinum beds in Tampa Bay also indicated the




Figure 3.5 Average density (short shoots m⁻²) of Halodule wrightii at control and three increasing levels of shading (30, 50, 80%) in Tampa Bay seagrass beds (± 1 S.D.). (Modified from Neely 2000)

maximum depth limit in the bay occurred at about 20.5% of subsurface PAR (Dixon and Leverone 1995, 1997). Another study on irradiance in the more oceanic waters at the mouth of Tampa Bay showed that, using 20% light penetration as the threshold, sufficient light was available for T. testudinum beds to extend to two meters of depth (Miller and McPherson 1995). That depth is similar to measured seagrass-bed depth limits, which ranged from 1.98 to 2.37 m in estuaries (Dixon and Leverone 1995). Evaluating short-shoot density, Neely (2000) found that shallow, monospecific beds of H. wrightii showed a direct relationship between density and irradiance level at Mullet Key in Tampa Bay (Figure 3.5). The same level of density reduction with shading occurred in the cool and warm seasons, although summer short-shoot densities were almost twice those of the winter densities, indicating the critical effect that underwater irradiance has on seagrasses. Similar declines in leaf growth, short-shoot density, biomass and primary production have been caused by shading T. testudinum (Lee and Dunton 1997; Kraemer and Hanisak 2000) and H. wrightii (Czerny and Dunton 1995).

Light attenuation that limits seagrass depth distribution can result from a variety of sources (Kenworthy and Haunert 1991; Morris and Tomasko 1993). One factor governing light effects on seagrass depth distribution is the level of epiphytic biomass. The average annual attenuation by epiphytes on water-column PAR levels available to *T. testudinum* blades was from 32%–47% in Tampa Bay, 40%–56% in Sarasota Bay, and 21%–44% in Charlotte Harbor (Dixon and Leverone 1997). Variation in epiphyte loads on *T. testudinum*

blades resulted in a 7%–67% seasonal range in PAR attenuation in Tampa Bay, 13%–99% in Sarasota Bay, and 6%-89% in Charlotte Harbor (Dixon and Leverone 1997). Thus, in determining seagrass light requirements, seasonal levels of epiphyte loads should be considered in addition to the actual PAR levels (Dixon 2002).

Other factors shown to have critical effects on light available to seagrass, in a light model developed for the Indian River Lagoon, were total suspended solids (e.g., tripton or non-algal particulate matter) and chlorophyll a concentration, which were the dominant causes of light attenuation affecting seagrass depth distribution (Christian and Sheng 2002). Attenuation of light by turbidity is a critical abiotic factor governing availability of light to seagrasses, as shown by the effects of dredging in the Laguna Madre of Texas that resulted in the decline of H. wrightii (Onuf 1994). Depth distribution, biomass, and photosynthetic responses were compared for shallow and deep populations of T. testudinum on the Gulf coast of Florida (Dawes and Tomasko 1988; Dawes 1998b). In general, biomass decreased while chlorophyll a increased with greater depth (Table 3.2), and the ratio of above- to below-ground biomass remained similar.

Also, light quality (i.e., wave-length distribution) and PAR affect seagrass growth. Exposure of H. wrightii, H. engelmannii, and S. filiforme to UV-A and PAR irradiation resulted in photosynthesis inhibition in the latter species (Trocine et al. 1982). A low ratio of red:far-red wave lengths reduced branching and node formation in R. maritima, a response that may allow plants shaded by larger plants to escape by emphasizing horizontal rhizome elongation (Rose and Durako 1994). A similar response by H. wrightii, when shaded by T. testudinum, was noted in the Florida Keys (Tomasko 1992).

Photosynthetic responses have been used to evaluate seagrass tolerances to salinity and temperature variations (Dawes *et al.* 1987, 1989). Estuarine populations of *H. engelmannii* and *H. johnsonii* from the Gulf coast and Indian River Lagoon of Florida, respectively, had broader tolerances to a range of temperatures and salinities when compared to Gulf-coast oceanic populations of *H. engelmannii* and *H. decipiens*. Differences in photosynthetic factors (*e.g.*, P_{max} , α , I_k) were evident with freshly collected plants and after culturing

Table 3.2 Photosynthetic characters (P_{max} , alpha, I_c , total chlorophyll), primary production, and leaf growth (\pm S.D.) of *Thalassia testudinum* in deep- and shallow-water populations at two Florida Gulf-coast sites.

Site	Leaf Relative Growth Rate (mg dw \cdot g dw ⁻¹ \cdot d ⁻¹)	Primary Production (g dw⋅m⁻²⋅d⁻¹)	Total Chl (mg⋅g dw ⁻¹)
Egmont Key			
Shallow (-0.4 m)	28.7 (5.4)	11.71	4.86 (0.99)
Deep (-2.5 m)	27.4 (6.3)	8.17	3.92 (1.28)
Anclote Key			
Shallow (-1.0 m)	31.1 (9.3)	1.39	3.00 (0.84)
Deep (-2.5 m)	20.5 (0.3)*	1.00	2.39 (0.70)
	P _{max}	I _c	Alpha
Site	$(\operatorname{mg} \operatorname{O}_2 \cdot \operatorname{g} \operatorname{dw}^{-1} \cdot \operatorname{h}^{-1})$	$(\mu E \cdot m^{-2} \cdot s^{-1})$	$(P_{\text{max}}/I_{\text{c}})$
Egmont Key		**	
Shallow	4.54 (0.70)	39	0.063
Deep	6.31 (1.21)	45	0.069
Anclote Key			
Shallow	6.08 (0.92)	32	0.103
Deep	2.60 (0.33)*	24	0.097

Modified from Dawes and Tomasko (1988). * = probability < 0.01

plants under common garden conditions for a number of weeks using various combinations of temperature and salinity. Salinity tolerance of *T. testudinum* from San Carlos Bay in the Caloosahatchee River Estuary in southwest Florida was assessed using long-shoot rhizomes bearing two to four short shoots (Doering and Chamberlain 2000). After a four-week acclimatization period in ambient seawater (35 ppt) in the laboratory, salinity was changed daily in 1.5 ppt increments.




Figure 3.6 Blade length and mean blade elongation rates for Thalassia testudinum cultured for 43 days in salinities ranging from 6 to 35 ppt. Letters indicate statistical differences between treatments and means, different letters are statistically different (P < 0.05). (Modified from Doering and Chamberlain 2000)

Regardless of other variables, blade length decreased (Figure 3.6) and blade production ended after 43 days in the lowest (6 ppt) salinity. Blade length was positively correlated with salinity after 43 days of exposure. Also, the number of blades per short shoot and blade biomass were similar in 12, 18, 25, and 35 ppt salinities after 43 days of exposure (data not shown).

Salinity affects the local distribution of seagrass species on the Gulf coast of Florida (Phillips 1960b, c), with euryhaline species such as H. wrightii able to grow in a broad range of salinities (e.g., 5 to 55 ppt; Dunton 1996). However, the effect on seagrass distribution due to salinity alone is difficult to separate from the effects of other factors, as shown in the general reduction of seagrass beds throughout Tampa Bay after the 1997–1998 El Niño event. The 1998 rainfall increase resulted in higher levels of dissolved inorganic nitrogen, chlorophyll a (phytoplankton), and color content (tannins) from extensive runoff into the bay (Johansson 2002a), which in turn reduced PAR penetration in the water column. In addition, the increase in rainfall directly lowered salinities, with Middle Tampa Bay salinities declining from a winter high of 30 ppt to 10 ppt by March and then to less than 5 ppt over a two-week period in April, 1998 (Dawes, unpublished data). Therefore, com-

Table 3.3 Structural features of *Ruppia maritima* epidermal cells from a single monoculture grown in 10 ppt and 32 ppt salinity.

Component of Cell Volume	Mean (± SD) at 10 ppt	Mean (± SD) at 32 ppt	Significance Level
Vacuolar volume 28.4% (± 19.8)		9.5% (± 9.3)	P = 0.03*
Cell membrane invagination	$0.5 \ \mu m \ (\pm \ 0.7)$	$5.0 \ \mu m \ (\pm 4.2)$	P = 0.001**
Polysaccharide volume	6.7% (± 3.5)	25.0% (± 13.4)	P = 0.03*
Chloroplast volume	8.6% (± 4.8)	$13.3\%~(\pm 5.1)$	P = 0.02*
Mitochondria number	12.2 (± 8.9)	17.3 (± 5.5)	$P = 0.03^*$

Modified from Kruzcynski (1994). * = significant; ** = highly significant.

bined with higher nutrient loadings and turbidity-caused light attenuation, lower salinities resulted in a die-back of *T. testudinum* in Cockroach Bay by June 1998 and a reduction in seagrass acreage throughout west-central Florida coastal areas (see Chapter 2).

Seagrass osmoregulatory responses to changing salinities involve structural and physiological changes. The epidermal cells of T. testudinum and R. maritima show cell-wall and organelle modifications when grown or collected in salinities above 30 ppt (Jagels and Barnabas 1989). In R. maritima, clones of a single genetic strain cultured in 32 ppt had convoluted cell membranes, smaller vacuoles, and numerous mitochondria associated with the cell membrane (Table 3.3). Also, plasmodesmata were not evident between the epidermal and cortical cells, and a significant increase occurred in acidic mucopolysaccharides associated with the inner cell wall (Kruzcynski 1994). Clones from the same genetic strain of R. maritima did not show any of the above features when cultured at a lower salinity (ca. 10 ppt). The increase in cell membrane surface area and associated mitochondria are characteristic of cells active in osmoregulation, and acidic carbohydrates may be involved in ion transport and binding (Dawes 1998a). In addition, osmotic roles of the amino acids proline, alanine, and glutamate have been identified in seagrasses (Pulich 1986; Adams and Bate 1994). Pulich (1986) reported that proline functions as an organic osmoticum in R. maritima, T. testudinum, and H. wrightii, and alanine serves that role in H. engelmannii.

Data on temperature effects on seagrasses are limited to studies examining flowering and seed germination (see earlier discussion) and from a study of leaf-blade production near an electricitygenerating plant (Barber and Behrens 1985). In the latter study, the productivity of *S. filiforme* increased in cooler months and decreased in warmer months in areas where water temperatures were enhanced by thermal pollution, whereas *T. testudinum* showed little or no response.


Water movement influences the spatial distribution of seagrass beds, in addition to the effects on depth distribution from tidal and wave activity (see also Chapter 5). Water velocities, degree of exposure to waves, and relative water depths strongly affect the distribution of H. wrightii and Z. marina beds near Beaufort, North Carolina (Fonseca and Bell 1998). Water movement, expressed as bladefriction velocities (u* = cm s⁻¹), reduced boundarylayer effects around the leaf, and photosynthetic rate increased in T. testudinum, up to a current velocity of 0.25 cm s-1 (Koch 2001). In addition, increased water movement causes a decrease in concentration of sediment sulfide, a phytotoxin. Further, seedlings of *T. testudinum* collected in the Florida Keys showed optimum growth under intermediate flow rates ($u = 0.3 \text{ cm s}^{-1}$), whereas stagnant water ($u \times = 0.0$) contributed to lower biomass possibly because of increased sulfide levels, and high flow rates ($u \times = 1.0 \text{ cm s}^{-1}$) reduced nutrient availability in the sediment pore water (Koch 1999a). Back-and-forth wave motion increases nutrient exchange between blades and the water column, in contrast to unidirectional movement or currents (skimming) that flow over T. testudinum meadows in the Florida Keys (Koch and Gust 1999). When the flow rate (u*) equaled 25 cm s⁻¹, the water current began to erode the sand; this caused a 50% reduction in seagrass coverage.

Repeated intertidal exposure, and thus the development of tolerance to desiccation, has been suggested as a causal factor in the ability of

H. wrightii to grow intertidally, where it may form extensive beds, in contrast to the subtidal distribution of T. testudinum (Phillips 1960a). However, factors other than desiccation tolerance may play a more critical role in terms of the higher vertical zonation for H. wrightii, for example, a greater tolerance to high irradiances and benefits from higher nutrient input from the shore (Björk et al. 1999).

Tolerance of exposure to high light levels in shallow or intertidal waters suggests that seagrasses may have evolved photosynthetic mechanisms (e.g., C4 photosynthesis) that reduce damage by UV radiation or intense PAR. However, photosynthetic mechanisms of all seagrass studied thus far appear to involve carbon-fixation reactions characteristic only of the Calvin cycle (C3 plants) (Beer et al. 1977, 1980; Andrews and Abel 1977; Benedict et al. 1980; Durako 1993), with the possible exception of the Mediterranean species, Cymodocea nodosa (Beer et al. 1980). Carbon fixation in C₃ plants occurs via the Calvin Cycle, in which the enzyme ribulose-1,5-bisphosphate carboxylaseoxygenase (rubisco) initially produces a three-carbon (C₃) compound, phosphoglycerate (Abel and Drew 1989; Beer 1996). However, an earlier study (Benedict and Scott 1976) reported that high levels of labeled carbon (14C-bicarbonate) were incorporated into malate (30%) and aspartate (33%) in T. testudinum. These two four-carbon acids are commonly associated with the initial fixation step of the Hatch-Slack pathway of C₄ plants. The process described by Benedict and Scott (1976) for T. tes-

Figure 3.7 Stable-carbon isotope values of seagrasses are influenced by adjacent mangroves.



tudinum is similar to Crassulacean Acid Metabolism (CAM), which prevents loss of fixed carbon (as CO₂) resulting from photorespiration suffered by plants under high light irradiance. Because seagrass lacunae store respired CO₂, determining the level of photorespiration in seagrass blades is difficult, although it appears to be lower than in susceptible terrestrial plants (Abel and Drew 1989). More studies are needed to determine whether any of the six Gulf-coast species have modified C₄ carbon-fixation pathways. Durako (1993) demonstrated that *T. testudinum* can utilize both bicarbonate (HCO₃-) and carbon dioxide, with relatively efficient HCO₃- utilization, despite an apparently low affinity for this form of carbon.

Plants with C₄ photosynthesis have higher (less negative) δ¹³C values than those of plants using only the C3 pathway. It has been shown that seagrasses have δ¹³C values similar to those of C₄ plants. The similarity in δ^{13} C values may be due to diffusional restrictions on carbon movement into the chloroplast in a relatively closed carbon-fixation system rather than reflecting their mode (e.g., C₃ or C₄) of carbon fixation (O'Leary 1988). Overall, average seagrass δ^{13} C values range between -10and -11 ppt, showing stable-isotope-ratio signatures that are usually less depleted in 13C than they are in other aquatic primary producers (Hemminga and Mateo 1996). Thus, δ13C values can be used to trace carbon flow in short-term carbonallocation and production studies. In contrast to the overall averages for seagrasses given by Hemminga and Mateo, the δ^{13} C values for T. testudinum leaves in south Florida, when growing adjacent to mangroves (Figure 3.7), ranged from -7.3 to -16.3 with stable-isotope ratios of carbon that were similar to those of mangrove detritus (Lin et al. 1991). The difference in δ^{13} C values suggests influence of mangrove detritus on nearby seagrass beds. Macroalgae may have nitrogen stable-isotope ratios similar to those of seagrasses, as shown for δ¹⁵N. Using δ¹⁵N values, Dillon et al. (2002) determined the sources of nitrogen in Florida. In Sarasota Bay, macroalgae had δ15N values that averaged +4 per mil greater than those of seagrasses, the higher values being positively correlated with δ15N values of wastewater discharge and being similar to macroalgae δ15N values near populated areas in the Florida Keys.

Seagrasses facilitate the transport of inorganic carbon from seawater to the chloroplasts of their

epidermal cells to make up for slow CO2 diffusion rates and low CO2 levels in water, as well as for the low affinity of rubisco for CO2 when the lacunae have high concentrations of O2 (Beer 1996). More importantly, seagrasses can use HCO₃- from seawater via carbonic anhydrase-catalyzed extracellular conversions. Unlike seaweeds, which have photosynthetic rates that are saturated by present-day CO2 levels, seagrasses continue to be CO₂-limited (Beer and Koch 1996). The authors suggest that when seagrasses colonized the sea in the Cretaceous, CO2 levels were higher. With atmospheric CO₂ levels again increasing, the authors suggest that nearshore beds may expand (but see Beer et al. 2003). However, CO2 concentration in seawater need not be a limiting factor to seagrass growth under low irradiance conditions, as shown by Durako and Hall (1992). Using δ¹³C values, they reported that CO₂ concentration in natural seawater was not limiting for T. testudinum under low light when photosynthetic rates were low, a relationship that may be relatively common in turbid water.

Entire-plant carbon budgets have been measured for *T. testudinum* in Florida Bay (Fourqurean and Zieman 1991) and for *H. wrightii* in Tampa Bay (Neely 2000). Below-ground biomass (roots, rhizomes, and short shoots) of *T. testudinum* accounted for 85% of the standing stock and 57.5% of total-plant respiration (Fourqurean and Zieman 1991). In contrast, below-ground biomass of *H. wrightii* ranged from 94% (winter) to 69% (summer) in Tampa Bay, with an above-ground productivity of 0.25 g C m⁻² d⁻¹ in September (Neely 2000).

Tomasko and Dunton (1995) compared four methods (blade clip-and-reharvest, above-ground biomass values, leaf-production turnover rates, and below-ground:above-ground-biomass ratios) for estimating diel rates of primary production in H. wrightii populations of Texas. The whole-plant biomass and leaf-turnover approaches appeared to be useful tools for estimating productivity in H. wrightii. The blade clip-and-reharvest method devised by Virnstein (1982) for determining primary production in narrow-bladed seagrasses such as H. wrightii, underestimated primary productivity. Further, they concluded that the H_{sat} model that calculates total PAR for a photoperiod was reliable and accurate in the prediction of daily carbon gain. However, they stated that although the in situ measurement of underwater light levels and P vs. I parameters, as used in Zimmerman *et al.* (1994), is a more rigorous approach for calculating primary production, it requires a large investment of time and resources.

Because of their conspicuous roles in productivity and biomass generation, both the organic composition and the caloric values of seagrasses have been studied (Dawes 1998a). Annual mean caloric values per gram of dry weight (g dry wt) for leaves, short shoots, and rhizomes of T. testudinum were 10-13, 8-11, and 12-14 kJ, respectively. Similar values were obtained for S. filiforme (10–13, 10–14, and 15–16 kJ, respectively) and H. wrightii (13-15, 12-13, and 14-16 kJ, respectively). Another study in seagrass beds between Tarpon Springs and Cedar Key estimated that available kilocalories ranged from a 16-month low of 344 to a high of 1,837 kcal⁻¹ m⁻², with the highest SAV biomass and caloric values occurring in the late spring and summer (Dawes et al. 1987).

Dawes (1998a) found that lipid levels for *T. testudinum*, *S. filiforme*, and *H. wrightii* were low, ranging from <0.1% to 6.3% of the dry weight for various organs. Soluble protein ranged from 5% to 22% of the dry weight, with leaves having the highest content in all three species. Soluble carbohydrates ranged from 6% to 54% of the dry weight, with rhizomes having the highest levels. Soluble protein and soluble carbohydrate levels show pronounced seasonal and species variations, as summarized by Zieman and Zieman (1989) and Dawes (1998a).

Rhizomes, followed by short shoots, are the principal storage organs for soluble carbohydrates and proteins in T. testudinum, S. filiforme, and H. wrightii (Dawes and Lawrence 1980) and in H. engelmannii (Dawes et al. 1987). Further, rhizomes are the source of soluble carbohydrates and proteins for regrowth of cropped blades and for initiation of spring blade growth in T. testudinum (Dawes and Lawrence 1979, 1980; Tomasko and Dawes 1989). In a review of carbon, nitrogen, and phosphorus content in the leaves of 27 seagrass species, Duarte (1990) found that they accounted for 34%, 2%, and 0.2% of the dry weight, respectively, with a mean C:N:P ratio of 474:24:1. Although the cell walls of seagrasses contain lignin, celluloses, and hemicelluloses, they are in lower concentrations than in terrestrial monocots (Table 3.4). Soluble, nonstructural carbohydrates (hemi-

Table 3.4 Cell-wall constituents of seagrass leaves, including lignin, cellulose, and soluble carbohydrates.

Species	Site (Ref.)	Lignin	Cellulose	Sol. Carb.
Thalassia testudinum	Florida (1)	0.2-2%	18–32%	1–9%
Syringodium filiforme	Florida (1)	1-3%	19-26%	10-16%
Halodule wrightii	Florida (1)	2-4%	25-33%	4-14%
H. uninervis	Gulf of Acaba (2)	<1%	41%	_
Halophila stipulacea	Gulf of Acaba (2)	2%	36%	_
H. ovalis	Gulf of Acaba (2)	<1%	33%	_
Heterozostera tasmanica	Philip Bay (3)	5%	20%	_

Modified from Dawes (1986). l. Dawes (1986); 2. Baydoun and Brett (1985); 3. Webster and Stone (1994).

celluloses) can account for 1%-16% of cell-wall dry weight in seagrass blades, whereas structural cellulose and lignin account for 18%-40% and 0.2%-5% respectively. By comparison, tropical grasses such as tall fescue (Festuca arundinacea Schreb.) and Bermudagrass (Cynodon dactylon (L.) Pers.) have 37% and 40% hemicellulose, 32% and 27% cellulose, and 10% and 6% lignin per g dry wt1, respectively (Dawes 1986). The adaptations of seagrasses to a hydrophytic environment are seen in their flexible blades (i.e., few fiber bundles) and low amounts of lignin. Perhaps the brief life spans of leaves of many seagrass species, ca. 15 days in T. testudinum (Witz and Dawes 1995; van Tussenbroek 1996b), may not allow time for significant production of lignin to occur, compared to the long leaf life spans (months) of terrestrial plants, thus making seagrass blades more edible to grazers.

RHIZOSPHERE and SEDIMENT DYNAMICS

Seagrasses serve as sediment traps by acting as baffles, causing sediment-laden water to slow and drop its sediment as shown for seagrasses of Florida Bay (Prager and Halley 1999). Even species with diminutive morphologies, such as *H. decipiens*, can reduce sediment movement (Fonseca 1989). However, studies in *T. testudinum* beds in Tampa Bay show that sediment resuspension does occur because of tidal currents, so these communities are both sinks and sources of suspended matter (Koch 1999b).

Seagrasses modify the anaerobic sediment in which they grow. Roots and rhizomes release oxygen translocated from the shoot, cause bioturbation of the soil, and add organic matter via decomposition (Moriarty and Boon 1989). Chem-

ical changes wrought by the below-ground components are critical for plant survival because high levels of sulfides in anoxic reducing sediments are toxic to plants (Erskine and Koch 2000; Koch and Erskine 2001; Carlson et al. 2002). The above-cited experiments created sediment sulfide concentrations up to 13.0 mM; natural sulfide levels of healthy seagrass beds in Florida Bay average <2 mM, whereas in areas of seagrass die-off, the phytotoxin averages 4-10 mM (Carlson et al. 2002). Some evidence exists that die-off of overdeveloped beds of T. testudinum in Florida Bay may be due partly to lowering of plant resistance to the parasite Labyrinthula sp. during periods of increased environmental stresses such as high salinity, high or low temperatures, hypoxia, and high sediment sulfides (Durako 1994; Blakesley et al. 2002; see Chapter 5). Evidence for sulfide toxicity was found when sulfide addition (6 mM) in tank cultures was combined with high salinity and temperature over a period longer than 28 days (Koch and Erskine 2001), thus mimicking die-off conditions in Florida Bay. Carlson et al. 2002 suggested that the higher sensitivity of T. testudinum to sulfide levels than that of H. wrightii may be due to the former species having a higher below- to above-ground biomass ratio. They also reported high sedimentsulfide levels in other estuaries, including Sarasota Bay, which may help explain the patchy distribution of seagrass beds (Figure 3.8). This suggests that carbonate sediments, as found in Florida Bay, are not the only type of substrate that contain high sulfide concentrations and that seagrass losses (but not die-off) in other areas on Florida's Gulf coast may also reflect sediment conditions. The release of oxygen into its rhizosphere by T. testudinum and the resultant decline in sulfides (converted to sulfates) was shown in seagrass beds in Laguna Madre, Texas (Lee and Dunton 2000a).




Figure 3.8 Frequency distribution of pore-water sulfide concentrations in carbonate sediments of Florida Bay (Rabbit Key Basin) and in siliceous, terrigenous sediments of Sarasota Bay (Carlson et al. 2002).

Much of the sulfide in anoxic sediments derives from the activities of sulfate-reducing microorganisms. Many microbial associations in the bulk soil and with the plant's roots can provide a means for obtaining scarce nutrients. For example, fungal associations (mycorrhizal) might provide phosphorus or micronutrients. Nielsen et al. (1999) hypothesized that the nature of seagrass sediments restricts the development of vesicular-arbuscular mycorrhizae (VAM) because they could not find the association in the rhizosphere of *T. testudinum*, in contrast to their finding that VAM are associated with submergent and emergent freshwater angiosperms. Because they did not find any (VAM) associations with the roots of T. testudinum, Nielson et al. (1999) suggested that formation of the association was restricted by the combined effects of highly anaerobic nature and high salinity of seagrass sediments.

Since the classic studies of Patriquin and Knowles (1972), who used acetylene reduction by the seagrass rhizosphere to infer nitrogen fixation, a number of studies have described nitrogen fixation and nutrient uptake by marine angiosperms in greater detail. Based on acetylene-reduction experiments, Kenworthy *et al.* (1987) determined that bacteria associated with the roots and rhizomes of *T. testudinum* fixed nitrogen. Nitrogen

fixation occurs in *Lyngbya wollei* (Phlips *et al.* 1992), a freshwater cyanobacterium. Perhaps *L. majescula*, a common mat and epiphytic species in seagrass beds of Florida's Gulf coast, has a similar capability.

Seagrasses take up nutrients principally from sediment pore water (e.g., ammonium and phosphorus; Fourqurean et al. 1992) but also from the water column (e.g., nitrate and ammonium; Touchette and Burkholder 2000). However, porewater nutrient concentrations in seagrass beds can vary widely, as shown in Florida Bay, where median values of 0.34 µM for soluble reactive phosphorus and 78.6 µM for ammonium occur (Fourgurean et al. 1992). Because of rapid growth rates, seagrasses require large amounts of fixed nitrogen (10 to 450 mg N m-2 d-1), with ammonium being the preferred form (Moriarty and Boon 1989). By contrast, nitrate and nitrite concentrations are usually low (<5 µM) in anaerobic sediments, presumably because they are rapidly used by denitrifying and anaerobic bacteria.

Seagrasses show a variety of responses when nutrients become limiting. Phosphorus is usually the most limiting nutrient in seagrass sediments of Florida Bay, where calcium carbonate rapidly binds any free PO₄-3 (Powell et al. 1989; Short et al. 1990; Touchette and Burkholder 2000), and its levels have been correlated with seagrass-bed development (Fourgurean et al. 1992). In Florida Bay, phosphorus apparently controls the successional sequence between early-stage H. wrightii beds and late-stage T. testudinum (Fourqurean et al. 1995). The former species tolerates higher nutrients better than T. testudinum does and will replace it in areas of natural or anthropogenic eutrophication. Nevertheless, factors other than nutrient levels may contribute to the replacement of T. testudinum by H. wrightii. Studies in Florida Bay (Carlson et al. 2002) reported that higher sulfide levels in the root zone, resulting from enhanced seagrass nutrition, can act as a phytotoxin in sediments and limit T. testudinum. By contrast, ammonium deficiency limits growth of T. testudinum in Laguna Madre, Texas (Lee and Dunton 2000b) and of S. filiforme in the Indian River Lagoon (Short et al. 1993).



- → Biomass production of the three Florida seagrass species with larger statures is highly variable, with highest values reported for each in summer and early fall. Below-ground biomass accounts for 50%–90% of total plant biomass.
- → Thalassia testudinum produces 10–19 leaf blades per short shoot per year, with lower turnover rates in more northern latitudes. Determinations of leaf production rates for other species are needed.
- Although vegetative growth is the primary method for seagrass expansion, little is known about rhizome extension rates for most species.
- → Seagrass communities serve as habitats and nurseries for over 170 species of invertebrates and more than 100 species of fish. About 60 fish species using seagrass beds as habitat have noncommercial values and have poorly known requirements.
- Epiphytic biota are often seasonal, serve as food, and enhance seagrass community structure but may also negatively influence seagrass growth by attenuating light.
- → A positive relationship exists between epiphytic load on seagrass blades and nutrient enrichment of their environment, whether the source of the nutrients is natural (e.g., bird rookeries) or human (e.g., septic tanks or stormwater runoff).
- → Drift macroalgae deposited on seagrasses can result in seagrass die-back but can also serve as habitat, as nutrient sinks, as food, and as a transport vehicle for invertebrates and fish.
- → Unlike overgrazing, which has been documented twice in Florida waters, moderate grazing may enhance the productivity and expansion of *T. testudinum*. The role of secondary metabolites that may reduce grazing in seagrasses is not known.
- → Data regarding competition between seagrass species or with their associated macroalgae are limited.
- Animal species using seagrass habitats also often use nearby salt marshes, mangrove forests, and coral reefs.
- → The export of inshore seagrass detritus offshore into deep water is poorly documented for Florida waters but may be an important source of organic matter in deep marine systems.

Manateegrass, Syringodium filiforme, with drift algae

FWC-FWRI photo

SEAGRASS COMMUNITY ECOLOGY

Like Florida's coastal mangrove and salt-marsh communities, seagrass communities are important primary producers, stabilize mobile sediments, serve as habitats and nurseries, and are direct and indirect food for diverse fauna (Dawes 1998a). Further, the characteristics of seagrasses (e.g. short-shoot densities, presence of inflorescence scars, levels of storage products) can aid in determining whether seagrass beds are stressed, thereby contributing to the monitoring of conditions in coastal and estuarine communities (Dennison et al. 1993; Dawes 1998a).

In a study of seagrass distributions and conditions on the northeastern Gulf of Mexico, Livingston et al. (1998) listed four significant predictors of seagrass community health. These predictors are photic depth, light quality, water quality (color, dissolved organic carbon, and chlorophyll a), and sediment characteristics. The principal predictor is light quantity (duration and intensity) in relation to seagrass depth distribution, which is related to water transparency, as shown in the review on seagrass depth limits by Duarte (1991). In this regard, water transparency was found to be the principal factor influencing the depth distributions of T. testudinum in Tampa Bay (Dixon and Leverone 1995, 1997) and growth form of H. wrightii below a canopy of T. testudinum in the Florida Keys (Tomasko 1992). However, light quality influences seagrass morphology as shown by altering the red to far-red ratios in cultures of Ruppia maritima (Rose and Durako 1994).

Seagrass communities have been measured for standing stock biomass (g dry wt m⁻²) and shortshoot densities (numbers m⁻²). Biomass data are commonly limited to above-ground material, mostly the photosynthetic blade tissue. Such data are available for the larger-stature seagrass species, including *T. testudinum*, *H. wrightii*, *R. maritima*, and *S. filiforme* (Table 4.1), but data are usually lacking for the smaller-stature species (*i.e.*, species of *Halophila*). In general, the below-ground biomass (roots and rhizomes) constitutes 50% to 90% of standing stock for the larger-stature species (Zieman and Zieman 1989). For example, total biomass estimates (above and below ground) for *T. testudinum* in Florida Bay ranged from 195 to

2,254 g dry wt m⁻², and 85% of this was below ground (Fourgurean and Zieman 1991). Beds of S. filiforme in the Indian River Lagoon have an annual mean above-ground leaf biomass of 87 g dry wt m⁻² (\pm 151) or 43% of the total, whereas below-ground roots and rhizomes accounted for 117 g dry wt m⁻² (\pm 94) or 57% of the total plant biomass (Short et al. 1993). Dunton (1990) found that in the Laguna Madre of Texas below-ground biomass for H. wrightii was 50% to 85% of the total and for R. maritima was 20% to 70% of the total biomass. On Florida's Gulf coast, the greatest leaf biomass for T. testudinum occurs in spring and early summer, whereas below-ground biomass (roots and rhizomes) is greatest in the fall (Dawes et al. 1985). Unlike below-ground biomass, leaf biomass in T. testudinum can show a four-fold seasonal fluctuation, with above- to below-ground ratios of 0.17 to 0.4 in December and 0.4 to 1.0 in April in Tampa Bay (Dixon and Leverone 1995).

Leaf turnover rates were reviewed by Short and Duarte (2001), and techniques to measure leaf growth were compared by Tomasko and Dunton (1995) and van Tussenbroek (1996b). Leaf production (g dry wt shoot - y -) in T. testudinum was measured by puncturing leaf clusters at their bases and measuring new leaf production after a few days (Tomasko and Dawes 1989). In a review of leaf turnover rates for T. testudinum, Marbá et al. (1994) reported that 14 to 19 leaves are produced annually by each short shoot. Leaf longevity on short shoots of T. testudinum in the Tampa Bay area using leaf-scar chronology was 14.6 days at Sunset Park in Tarpon Springs and 19.2 days in south Tampa Bay (Witz and Dawes 1995) and 24 days in a lagoon in the Yucatán, Mexico (van Tussenbroek 1996b). Leaf production generally ranged between 10 and 19 leaves short shoot y in Florida Bay (Zieman et al. 1989), the Indian River Lagoon (Gacia 1999), Tampa Bay (Witz and Dawes 1995), Charlotte Harbor (Tomasko and Hall 1999), Lower Laguna Madre, Texas (Kaldy et al. 1999), and the Caribbean coast of Mexico (Marbá et al. 1994; van Tussenbroek 1996b). Populations growing in cooler waters had lower annual rates of leaf production (10-15 leaves shoot1) for the Indian River Lagoon and the Lower Laguna Madre in Texas compared to those in Tampa Bay. Determination of leaf turnover in narrow-bladed seagrasses, such as H. wrightii (Tomasko and Dun-

Table 4.1 Standing stock (biomass) and productivity of southeastern United States seagrasses and closely related species (congeners) from other areas. Procedures differ greatly and influence the ranges in biomass and productivity.

		Biomass	Productivity
Species	Locality	(g dry wt⋅m ⁻²)	$(g C \cdot m^{-2} \cdot d^{-1})$
Halophila			
H. ovalis	India	48	
	Western Australia	40-60	
H. engelmannii	Texas (Gulf coast)	1.6	
Thalassia			
T. testudinum	Florida (general)	20-8,100	
	Florida (Atlantic coast)	20-1,800	0.9–16.0
	Florida (Gulf coast)	75–8,100	
	SW Florida Bay (Gulf coast)	60-125	
	Indian River, FL (Atlantic coast)	890	
¥	Corpus Christi, TX (Gulf coast)	454-885	
	Mexico, lagoon (Gulf coast)	420-1,418	
	Mexico, reef (Gulf coast)	573-811	
	Cuba (Caribbean Sea)	20-800	9.3-12.5
T. hemprichii	Queensland, Australia	70	
Syringodium	-		
S. filiforme	Florida (general)	15-200	
	Indian River, FL (Atlantic coast)	27-81	
S. isoetifolium	New Guinea (East Indies)	327	
Halodule	•		
H. wrightii	Mullet Key, FL (Gulf coast)	120	0.25
8	Florida (general)	0.8	
	North Carolina (Atlantic coast)	105-200	0.5-2.0
	Laguna Madre, TX (Gulf coast)	70-300	
	Mississippi Sound (Gulf coast)	256	
H. uninervis	New Guinea (East Indies)	150	
Ruppia maritima	Texas (Gulf coast)	0-200	

Modified from McRoy and McMillan (1977), Zieman and Wetzel (1980), and Hillman et al. (1989).

ton 1995) and *S. filiforme* (Fry and Virnstein 1988), requires using an approach different from the leaf-punch technique. Leaf productivity by *S. filiforme*, determined using leaf clipping and photography, was 1.8 g dry wt m⁻²d⁻¹ (Virnstein 1982; see Short and Duarte 2001 for a detailed review of techniques).

Little is known about rhizome growth rates of Florida's seagrasses, yet this is the sole method for vegetative expansion. Reviews indicate that vegetative growth differs among species, rhizome elongation rates being higher for smaller-stature seagrass species (Marbá and Duarte 1998). In their review, mean rhizome elongation rates were highest for *H. wrightii*, the smallest stature of the larger Caribbean seagrasses (223, range; 81–365 cm yr⁻¹), followed by *S. filiforme* (123, range:

52-182 cm yr⁻¹) and then the largest stature species T. testudinum (69, range: 22–152 cm yr⁻¹). Short et al. (1993) found that S. filiforme rhizome elongation exceeded 100 cm y-1 in the Indian River. At Mullet Key in Tampa Bay, rhizome internode length and increase in seagrass patch size for H. wrightii were positively correlated; the larger the internodes of H. wrightii, the greater the increase in seagrass coverage at the edges of patches (Jensen and Bell 2001). The authors also found that a 100-fold increase in phosphorus, but not nitrogen, resulted in a significant increase in rhizome internode length. Jensen and Bell (2001) suggested that the observed patch-size patterns in seagrass beds (e.g. larger landscape features) may be determined by small-scale factors (e.g. nutrient availability).

Table 4.2 Densities of macroinvertebrates at various sites in Florida, arranged geographically from the Atlantic coast to the northern Gulf of Mexico. Single-density values are means; ranges generally represent several sampling sites. Several values are derived indirectly.

	Seagrass	Faunal	Mesh Size	Density	Number	Deferre
Study Site	Species*	Group	(mm)	(indiv. m-2)	Species	Reference
Card Sound	Tha	total	0.8	1,085	_	Brook (1977)
Bahia Honda	Tha	Amphipoda	1.0	102	5	Nelson (1980)
Florida Bay	Tha	total	0.8	2,794	52	Brook (1978)
Rookery Bay	Hal	Amphipoda	1.0	910	3	Nelson (1980)
Tampa Bay	Hal	Polychaeta	0.62	13,313	44	Santos and Simon (1974)
Secretaria Constitution (Constitution Constitution Consti	Tha	Polychaeta	0.62	33,485		Santos and Simon (1974)
Anclote Anchorage	Syr	total	1.0	2,347	68	Mahadevan and Patton (1979)
	Syr	total	0.5	9,538	58	Mahadevan and Patton (1979)
	Tha	total	1.0	3,724	63	Mahadevan and Patton (1979)
	Tha	total	0.5	18,916	50	Mahadevan and Patton (1979)
Apalachee Bay	Tha/Syr	total	1.0	1,782 to 2,424	72	Lewis and Stoner (1981)
-	Tha/Syr	total	0.5	3,154 to 4,754	86	Lewis and Stoner (1981)
	Tha	total	0.5	16,108	80	Lewis and Stoner (1983)
	Tha/Hal	Crustacea	0.5	6,716	46	Lewis (1984)
	Tha/Syr/Hph	total	0.5	2,827	170	Stoner (1980b)
Apalachicola Bay	Hal	total	0.5	38,780	58	Sheridan and Livingston (1983)
St. Andrew Bay	Tha	total	0.7	7,567	89	Saloman et al. (1982)
,	Hal	total	0.7	3,370	67	Saloman et al. (1982)
	Tha/Hal	total	0.7	4,192	86	Saloman et al. (1982)
Santa Rosa Sound	Tha	total	0.5	13,260	144	Morton et al. (1986)
	Hal, core	total	0.5	9,020	41	Stoner et al. (1983)
	Hal, suction	total	0.5	6,077	37	Stoner et al. (1983)

Modified from Virnstein 1987. *Hal = Halodule wrightii, Tha = Thalassia testudinum, Syr = Syringodium filiforme, Hph = Halophila engelmannii.

STRUCTURE, SHELTER, AND PREDATION

Seagrass beds function both as nurseries (Lewis and Stoner 1981; Lewis 1984; Virnstein 1987; Zieman and Zieman 1989; Fonseca et al. 1996b; Bell et al. 2001; Heck et al. 2003) and as habitats for invertebrates (Greening and Livingston 1982; Virnstein 1987; Valentine and Heck 1993), fish (Stoner 1983; Gilmore 1987), sea turtles (Williams 1988a), and manatees (Lefebvre et al. 1989). The role of seagrass meadows as nurseries was reviewed by Heck et al. (2003) who reported a strong link between seagrass abundance and abundances of juvenile fin fish and shell fish. The principal factor in this positive correlation was the structure that seagrasses added to the habitat.

Invertebrates add structure and serve as food, thus supporting increased diversity of other forms. For example, high densities of the mussel Modiolus americanus in St. Joseph Bay reached 2000 individuals m⁻², which enhanced invertebrate diversity and secondary production (Valentine and Heck 1993). The high variation of animal diversities among seagrass sites may be due in part to wave and current activity (Bell et al. 1994). The authors found that the mean numbers of fish and copepods and mean numbers of fish species were higher in low-energy seagrass beds (e.g., minimum exposure to wind direction, speed, and effective fetch in Tampa Bay and Onslow Bay, North Carolina). Densities of macroinvertebrates in seagrass beds on Florida's Gulf coast vary widely among sites (Table 4.2), ranging from 910 individuals m⁻² (Rookery Bay) to 33,485 individuals m-2 (Tampa

Bay). Invertebrate communities of seagrass beds in Florida are highly regional, and overlap occurs between subtropical, tropical, and warm-temperate faunas (Virnstein 1987). Holmquist et al. (1989) found that seagrass beds in Florida Bay play a secondary role as habitat for decapods and stomatopods compared with their reactions to winter cold fronts as a function of their positions on mud banks. Decapod crustaceans, such as caridean shrimps, usually dominate the larger fauna numerically, similar species of shrimp being widespread in Florida's Gulf-coast seagrass beds (Holmquist et al. 1989). The macrofauna varies widely in species composition and animal density over distance and time, including those of amphipods, gastropod molluscs, and polychaete worms (Virnstein 1987).

Because infaunal, epibenthic, and epiphytic animals of seagrass beds are prey for larger animals, fish are abundant there (Gilmore 1987). Principally, the larger-sized fish (e.g., spotted seatrout, Cynoscion nebulosus, and pinfish, Lagodon rhomboides) have been studied. Smaller nonfishery species (e.g., gobiids, syngnathids), which account for most of the resident species in seagrass beds, are less well known but include over 100 species (Gilmore 1987). As with invertebrates, fish faunas vary with geographic regions on Florida's Gulf coast and include warm-temperate species, eurythermal tropical species, and stenothermal tropical species (Gilmore 1987). Four subenvironments in Florida Bay had an overall average density of 11 fish m⁻² and a total of 56 species (Sogard et al. 1987). The two-year study found seagrass standing crop and litter to be the most important determinates of fish numbers.

Monitoring animal densities and diversity in planted seagrass beds is a useful method for determining habitat restoration success. A study of a three-year-old planted seagrass bed in Tampa Bay (Fonseca et al. 1996a, 1996b) found that shrimp, fish, and invertebrate densities were equivalent to those of natural communities, although shortshoot density was only one-third those of natural beds. Another study in Tampa Bay found that higher numbers of polychaetes occurred in two-year-old planted sites than in natural *H. wrightii* beds (Bell et al. 1993). The rapid increase of animals in planted beds suggests that the threshold value of habitat structure may be much lower than previously thought and that the lack of correlation

between patch size and animal abundance suggests other factors are involved in maintaining animal populations. Clearly, more studies are needed to determine animal population relationships in restored, created, and patchy seagrass beds.

Studies on seasonal animal-population fluctuations and patch formation in seagrass meadows suggest that they are not static, even under "stable" environmental conditions. For example, Tampa Bay seagrass beds experiencing more rapid water movement have a lower number of fish and harpacticoid copepods (Bell et al. 2001), which may also be influenced by organic matter content in the sediment. In St. Croix, Virgin Islands, tropical seagrass beds fluctuate in response to seasonal conditions (Williams 1988b). In addition to seasonal changes, severe plant die-off has occurred in Florida Bay, probably hastened by spatial variations in environmental stresses (Durako 1994; Zieman et al. 1999; see also Chapter 5). Water-column nutrient increases resulting in eutrophication have been cited as important environmental stressors that can shift species dominance in seagrass beds and influence sulfide production (Carlson et al. 2002). Williams (1987) found that T. testudinum could shade and replace S. filiforme beds in St. Croix. She also reported that the successional sequence from bare sediment began with rhizophytic (psammophytic) algae, then proceeded to S. filiforme, and ended with T. testudinum, as sediment nutrient concentrations increased over time (Williams 1990). In this regard, considering nutrient increases emanating from bird perches installed in Florida Bay, Fourquean et al. (1995) proposed that H. wrightii might replace T. testudinum in Florida Bay as sediment nutrient levels rise. However, this study did not consider the effect of increased sulfide levels in the sediment (Carlson et al. 2002) or the effect of phosphate and nitrate on seagrass growth (Powell et al. 1989).

The competition for space or the "space race" is evident by the numbers of macroalgae, cyanobacteria and microalgae (Hall 1988; Harris 1997) that grow on seagrass blades and that show strong seasonal changes. At Egmont Key in Tampa Bay, the five species of cyanobacteria and 20 species of epiphytic macroalgae varied in dominance (by algal division) during a 15-month study of *T. testudinum*. Brown algae dominated during the winter and spring months and cyanobacteria in the




Figure 4.1 Seasonal variations in the mean numbers of cyanobacteria and macroalgae, grouped by divisions, epiphytic on Thalassia testudinum blades at Egmont Key in Tampa Bay over a 15-month period (Hall 1988).

summer months (Figure 4.1). Hall (1988) divided the algae into those that (1) are characteristically epiphytic on seagrasses; (2) settle and reproduce on seagrass blades, but are commonly epilithic; and (3) germinate on blades and then break free before reproduction. A study of *T. testudinum* beds near Leffis Key and Siesta Key in Sarasota Bay (Table 4.3) found many species of centric and pennate diatoms, six species of cyanobacteria, and 21 species of macroalgae (Harris 1997). Algae epiphytic on H. wrightii and T. testudinum in Tampa Bay, Sarasota Bay, and Charlotte Harbor (Dixon and Leverone 1995; Dixon and Kirkpatrick 1995, 1999) and Santa Rosa Sound in Pensacola Bay (Macauley et al. 1988) change seasonally, species diversity being negatively correlated with water temperature. Gacia et al. (1999) found that sheetlike and filamentous species of green-algae genera (Enteromorpha, Cladophora) were less resistant to grazing fish than were species of more coarsely branched red-algae genera (Hypnea, Chondria, Acanthophora) in the Indian River Lagoon. Algae

epiphytic on seagrass blades can account for 46% of the total biomass and 60% of the benthic primary production, as found using stable-isotope analysis, in *H. wrightii* beds in Mississippi Sound (Moncreiff and Sullivan 2001).

Studies have shown that epiphytes, in combination with PAR attenuation and seasonal lowgrowth periods, depress blade growth and primary production and reduce the depth of T. testudinum in Tampa Bay (Dixon and Leverone 1995, 1997), Florida Bay (Frankovich and Fourgurean 1997), the Florida Keys and Keys in Belize, Central America (Tomasko and Lapointe 1991), and Perdido Key in northwest Florida (Wear et al. 1999). Tomasko and Lapointe (1991) found water-column dissolved inorganic nitrogen (DIN) concentration was high near Big Pine Key (3.96 µM), an island in the Florida Keys with over 2,000 septic tanks. This DIN level is similar to that of a mangrove island in Belize (Man-O-War Key) having 75 pairs of nesting frigate birds (3.80 µM). In contrast, DIN levels were low around Cutoe Key (0.50 μM), a remote mangrove island in the Florida Keys. The seagrass beds near Big Pine Key and Man-O-War Key had higher epiphytism, lower short-shoot densities, lower leaf-area indices, and lower biomass in contrast to those near Cutoe Key (Tomasko and Lapointe 1991). Wear et al. (1999) reported similar results from an experiment using slow-release fertilizer (Osmocote®) in northwestern Florida at Perdido Key. Nutrification caused significant increases of diatoms, as well as red and brown macroalgae, on blades of T. testudinum, H. wrightii, and S. filiforme over a 12-month period. Frankovich and Fourgurean (1997) reported that epiphytic loads on T. testudinum, along a nutrientenrichment gradient in Florida Bay, gave similar results to the previous studies, and this study indicated that the effect is localized. The authors concluded that epiphyte levels alone are not as responsive to moderate nutrient enrichment as are other seagrass characters, such as leaf tissue N:C ratios. As noted in Chapter 3, epiphytes of seagrasses reduce water-column PAR by as much as 99% seasonally. The question of whether organic carbon, nitrogen, or phosphorus can be transferred from seagrass blades to the epiphytic community or vice versa is not clear (see Lobban and Harrison 1994).

Some of the most abundant invertebrates in

Table 4.3 Epiphytic algae on blades of *Thalassia testudinum* at Leffis Key and Siesta Key in Sarasota Bay (modified from Harris 1997) and at Egmont Key in the mouth of Tampa Bay (modified from Hall 1988).

Species	Leffis Key	Siesta Key	Egmont Key
Cyanobacteria			
Anabaena oscillarioides	x	X	x
Anacystis sp.	X	X	x
Calothrix crustacea	X	X	\mathbf{x}
Entophysalis conferta	X		
Microcoleus lyngbyaceus	x	X	x
Porphyrosiphon notarisii	X		
Chlorophyta			
Boodleopsis pusilla	x		
Chaetomorpha minima		X	X
Cladophora spp.	X	X	X
Enteromorpha chaetomorphoides			X
Ulva lactuca	x		
Phaeophyta			
Cladosiphon occidentalis	x		X
Ectocarpus elachistaeformis			x
Ectocarpus rallsiae (as Giffordia rallsiae)	x	X	X
Ectocarpus sp. (as E. rhodochortonoides)	x	x	X
Hincksia mitchellae (as Giffordia mitchellae)	x	x	X
Hummia onusta	x	x	X
Myrionema magnusii (as M. orbiculare)	x	X	X
Myriotrichia occidentalis	x	X	X
Rhodophyta			
Acrochaetium spp.	x	X	X
Centroceras clavatum	X	X	X
Ceramium cimbricum (as C. fastigiatum)	x	X	X
C. flaccidum (as C. byssoideum)	X	X	x
Chondria dasyphylla	X	X	x
Chondrophycus papillosus (as Laurencia papillosa)	x		
Erythrotrichia carnea			x
Hydrolithon farinosum (as Fosliella farinosa)	x	X	x
Hypnea musciformis	x	X	
Polysiphonia subtilissima	x	X	
Stylonema alsidii (as Goniotrichum alsidii)			x

seagrass beds are the meiofauna, including crustaceans (Lewis 1987) and copepods (Hall 1988). Harpacticoid copepods are often the most common animal group associated with seagrasses (Hall and Bell 1988, 1993; Walters and Bell 1986; Walters 1991). One study reported that more than 20,000 harpacticoid copepods migrated h⁻¹ m⁻² from the sediment during the night into the water column, representing over 50% of the total benthic harpacticoid community (Walters and Bell 1986). Further, this postsunset entry into the water column from the sediment involved over 30 species in 15 families (Walters 1991). At Mullet Key in

Tampa Bay, artificial blades (green ribbon) with artificial epiphytes (cotton fibers) attached had significantly higher densities, compared to green ribbons alone, of copepods, polychaetes, and nematodes associated with them after three days. This shows that epiphytic biota enhance structural complexity of seagrass blades, thereby increasing habitat values (Hall and Bell 1988). In a similar study, Hall and Bell (1993) showed that the most abundant meiofaunal groups on seagrass blades at Egmont Key were adult harpacticoid copepods, copepod nauplii, and nematodes. They also determined that meiofauna density was most signifi-




Figure 4.2 Drift algae are entangled by seagrass leaves, as in this Thalassia testudinum bed.

cantly correlated with percentage cover of filamentous algae. A 15-month study of *T. testudinum* blades at Egmont Key at the mouth of Tampa Bay (Hall 1988) showed seasonal shifts in the abundance of nematodes (highest in winter) but not that of harpacticoid copepods and nauplii (abundant throughout the year). Epiphytes trap detritus, which attracts significantly higher numbers of copepods that feed on it (Meyer and Bell 1989). Small grazers (e.g., copepods) can play a role in epiphyte control (Virnstein 1987), as shown for *H. wrightii* plants in the Indian River Lagoon (Howard and Short 1986).

Along with drift macroalgae, rhizophytic (psammophytic) algae may equal or surpass seagrass biomass in Florida Gulf-coast seagrass beds (Dawes et al. 1985). With the exception of some tropical coenocytic green algae, macroalgae usually cannot attach to sand or mud. Coenocytic green algae are a diverse group of siphonaceous (lacking internal cell walls) algae placed in the order Caulerpales. They form extensive rhizoid holdfasts and include species of calcified (Halimeda, Penicillus, Udotea) and noncalcified (Caulerpa, Riphilia, Avrainvillea) genera. Even though rhizophytic coenocytic algae may compete for space and light with seagrasses, they may also serve as pioneer plants, which then may be replaced by seagrasses, as shown in a coral-reef lagoon in St. Croix (Williams 1990).

Over 230 species of unattached macroalgae (drift algae) are known throughout the world

(Norton and Mathieson 1983). Drift algae are usually noncalcified, remain infertile, and reproduce vegetatively (Collado-Vides et al. 1994). Two studies, one in Tampa Bay, Boca Ciega Bay, and Anclote Anchorage (Phillips 1960b) and the second in Crystal Bay (Phillips 1960c), listed 195 and 46 taxa, respectively, of algae that could not be distinguished as either drift or attached. Drift algae are concentrated in seagrass beds where water-current velocities are reduced and blades will entangle the algae (Figure 4.2). To some extent, seagrass blade size may influence entanglement (Bell et al. 1995). Seagrass beds on the Gulf coast of Florida are known to contain from 8 to 65 species of drift algae with the total biomass ranging from 0.002 g to 930 g dry wt m⁻² (Table 4.4). Brown (2001) found that algal biomass differed for three sites on the Gulf coast of Florida, both by season and by site. Annual means of 6.01, 3.83 g, and 2.25 g dry wt m⁻² were obtained for Tampa Bay in Cockroach Bay, Tarpon Bay, and Sunset Beach respectively. Although biomass varies with season and site on the Gulf coast, the types and numbers of species are similar (Table 4.4).

Drift algae are an important ecological component of seagrass beds because they serve as habitat, provide transportation, and provide a food source for invertebrates (Ballantine *et al.* 1994; Holmquist 1994; Greenway 1995; Knowles and Bell 1998; Maciá 1999; Brooks and Bell 2001; Rydene and Matheson 2003). In Florida Bay, Holmquist (1994) found 61 invertebrate taxa in drift algae mats in

Table 4.4 Number of drift-algae species and dominant species, biomass, and presence/absence (by division) in seagrass beds on the Gulf coast of Florida. Locations include (1) Anclote Estuary near Tarpon Springs (Hamm and Humm 1976), (2) Hillsborough Bay including studies 2a (Kelly 1995) and 2b (Avery 1997), (3) Sunset Beach at Tarpon Springs (Brown 2001), (4) Cockroach Bay in Tampa Bay (Brown 2001), and (4) Tarpon Bay at Sanibel Island (Brown 2001).

	Locations					
	1	2a	2b	3	4	5
Number of species	65	19	18	9	9	8
Number of dominant species	4	5	9	9	9	8
Mean biomass (g dry wt ⋅ m ⁻²) low	-	_	80	0.07	0.0002	0.14
high	-	_	930	6.34	9.45	9.32
Cyanophyta						
Lyngbya majescula					X	X
Chlorophyta						
Caulerpa prolifera			X			
Chaetomorpha crassa			x			
C. gracilis			X			
Chaetomorpha sp.		X				
Enteromorpha spp.						x
Enteromorpha clathrata			x			
Ulva lactuca		X	X		x	
Phaeophyta						
Sargassum pteropleuron				x		
Sargassum spp.	\mathbf{x}					
Rhodophyta						
Acanthophora spicifera			x		X	X
Chondrophycus papillosus (as Laurencia papillosa)	X					
Digenia simplex	\mathbf{x}					
Gracilaria armata					X	X
G. caudata				x	X	X
G. cornea (as G. debilis)					X	
Gracilaria spp.		X	X			
Hypnea musciformis				x	X	X
Laurencia intricata (as Laurencia obtusa)	\mathbf{x}			\mathbf{x}		
L. poiteaui				x		
Polysiphonia ramentacea				\mathbf{x}		
P. subtilissima			X			
Solieria filiformis (as Agardhiella tenera)		\mathbf{x}	x	X	X	X
Spyridia filamentosa		X	X	x	x	X

seagrass beds. Average drift rate was 0.5 km d⁻¹, indicating that drift algae are an effective mode of transport for animals. Ballantine *et al.* (1994) reported that balls of algae in Venezuela contained 12 species of invertebrates. The numbers of amphipods, isopods, and tanaeids in Venezuela were significantly higher on two species of red algae than were associated with blades of *S. filiforme*, as similarly found in other previous studies (Knowles and Bell 1998). Virnstein and Howard (1987) noted that drift algae serve as habitat and food sources for gammaridean amphipods. Brooks

and Bell (2001) found that drift clumps of the red alga *Hypnea cervicornis*, as they moved through seagrasses and into sandy areas, had significantly higher amphipod densities; they concluded that drift algae clumps provide a mobile corridor for animals. In Tampa Bay, drift algae form an important alternative habitat for 12 of the 20 most abundant juvenile and small adult fish (Rydene and Matheson 2003). However, drift algae deposited in seagrass beds can also affect these habitats by causing a seagrass die-back as a consequence of light and oxygen reduction (Norkko *et al.* 2000). A

combination of algae deposition and sea-urchin grazing increased the die-back of T. testudinum in Florida Bay (Maciá 1999). The amount of driftalgae biomass can also have a negative correlation with seagrass bed expansion if dissolved nutrients are high, as shown for Hillsborough Bay (Avery 1997). In a 10-year study, Avery (2000) found that with improved water quality the 18 species of dominant drift algae declined in average biomass from 164 g dry wt m-2 in 1987 to 0.0029 g dry wt m⁻² in 1998 in Tampa Bay. The green coenocytic alga Caulerpa prolifera had colonized Hillsborough Bay in 1986 with an initial coverage of almost 220 ha (543 acres) but had declined to zero in 1997. In contrast, H. wrightii coverage increased from 0.2 ha (0.5 acres) in 1986 to nearly 57 ha (141 acres) in 1998 (Avery 2000).

A few studies exist concerning competition between seagrasses or between seagrasses and attached macroalgae, with two reports on T. testudinum, S. filiforme, and H. wrightii and rhizophytic algae in the Caribbean (Williams 1987, 1990). Williams (1987) removed the leaf canopy of T. testudinum in St. Croix, resulting in increased irradiance reaching the S. filiforme understory, and its biomass doubled within nine months. A study in Tampa Bay indicated that competition occurred between T. testudinum and H. wrightii (Rose and Dawes 1999), where T. testudinum had a lower biomass when growing with H. wrightii, suggesting interspecific competition. Further, leaf biomass and growth were significantly lower in dense, monotypic beds of T. testudinum compared to less dense beds, suggesting light was reduced via shading in dense beds (Rose and Dawes 1999). There are also reports regarding competition between H. wrightii and the coenocytic green macroalga C. prolifera (Bottone and Mattson 1987; Bottone and Savercool 1993). Although the data are not conclusive, C. prolifera may act as a weak competitor for space with H. wrightii, coexisting with it as an opportunistic species under stable environmental conditions in Tampa Bay (Bottone and Savercool 1993). This result is similar to those found in studies in the Indian River Lagoon (White and Snodgrass 1990). Further, C. prolifera, although susceptible to sudden environmental changes (e.g., low salinity), can rapidly colonize areas devoid of vegetation that may occur following seagrass dieback (Bottone and Mattson 1987).

Overgrazing of seagrass beds is a rarity but has occurred on the Florida Gulf coast. Overgrazing by Lytechinus variegatus (purple sea urchin) resulted in large-scale denudation of seagrass beds. Over 20% of an area 26 km by up to 9.5 km were consumed near the Pepperfish Keys south of the Steinhatchee River in the northern peninsula (Camp et al. 1973). Density at the front of a single aggregate averaged 636 sea urchins m⁻², with individuals piled upon one another 2-8 individuals deep that covered the substrate. Dense aggregations of the same urchin species overgrazed about 0.81 km² of T. testudinum in outer Florida Bay; mean densities ranged from 364 to <1 individual m⁻² (Rose et al. 1999). The origins of the urchin feeding fronts are not understood. In Florida Bay, high urchin densities may reflect unusually high recruitment or a release from predation pressure because of overharvesting of spiny lobsters and stone crabs (Rose et al. 1999).

Moderate grazing pressures on T. testudinum blades were correlated with specific leaf growth rates (0.024 mg dry wt. d-1) in the Yucatán, Mexico; this showed that sporadic grazing had little detrimental effect on T. testudinum (Cebrián et al. 1998). Studies on T. testudinum beds in northwest Florida and a review of the literature indicated that herbivory plays a major role in stimulating seagrass growth by removing macroalgae (Heck and Valentine 1995; Valentine and Heck 1999). Also, long-term effects of grazers have been shown in a relationship between L. variegatus and T. testudinum in the Gulf of Mexico (Heck and Valentine 1995), where urchins and seagrass coexist in balance. In these cases, a balance developed among intensive grazing, loss of habitat, and predation by fish on the urchins. In the 1995 experiment, protection from predator fish by enclosures resulted in intense grazing by the urchins, which became most destructive in winter when T. testudinum could not recover as rapidly. The more intensively grazed beds showed significant reductions in above- and below-ground biomass, which was apparent even 3.5 years after grazing had ceased. Varying the number of L. variegatus in experimental plots in St. Joseph Bay, Valentine et al. (1997) found an increase in shoot density and productivity of T. testudinum under moderate grazing pressure. In a later study, Valentine et al. (2000) demonstrated that the effect of grazing in Florida Bay was related

Table 4.5 δ15N and δ13C values of seagrasses and marine macroalgae from various locations.

Plants	δ¹5N	δ ¹³ C
Seagrasses		
Thalassodendron ciliatum (Australia)	+3.5	-9.3
Syringodium isoetifolium (Australia)	+5.0	-4.0
Thalassia testudinum (Nicaragua)	+3.5	-13.2
Halodule wrightii (Texas)	+3.9	-10.3
Ruppia maritima (Texas)	+3.6	-7.9
Halophila engelmannii (Texas)	+3.8	-7.9
Thalassia testudinum (Texas)	+3.9	-8.9
Thalassia testudinum (Jamaica)	<u>+4.3</u>	<u>-11.1</u>
,	$\overline{x} = 3.9 \pm 0.5$	-9.1 ± 2.7
Marine Macroalgae		
Ulva fasciata (Texas)	+8.1	-14.6
Ulva lactuca (Texas)	+8.1	-14.6
Gelidium crinale (Texas)	+7.9	-14.3
Ascophyllum nodosum (Maine)	+8.0	-16.9
Fucus vesiculosis (Maine)	+8.1	-16.5
Acetabularia kilnori (Australia)	<u>+6.5</u>	<u>-9.5</u>
,	$\overline{x} = 7.8 \pm 0.6$	-14.4 ± 2.6

Macko (1981) cited in Fry et al. (1987).

to season and water depth, with grazing having a stronger effect in the winter or in seagrass beds at greater depths. Thus, repeating experiments in environments with differing physical conditions and in different seasons is critical in order to determine the effects of blade loss and seagrass bed maintenance.

Few studies have been carried out on predation within seagrass beds since the reviews by Orth *et al.* (1984) and Zieman and Zieman (1989). Based on feeding and caging studies, predation, especially by pinfish and pink shrimp, is probably a major biological interaction influencing invertebrate community structure (Virnstein 1987). In a study in St. Joseph Bay, increased biomass of a mussel (*Modiolus americanus*) in *T. testudinum* beds resulted in greater abundance, biomass, and annual production of other invertebrates (Valentine and Heck 1993).

TROPHIC DYNAMICS

The ultimate fate of plant biomass is mineralization to simple inorganic compounds, with a portion of seagrass leaf decomposition occurring while it is still attached to the plant. Nevertheless, most biomass decomposition occurs on and in the sediment of seagrass meadows (Hemminga and Duarte 2000).

A long-term study (419 days) in Laguna Madre, Texas of H. wrightii senescent blades suspended in the water column revealed that decomposition resulted in a loss of 36% of the organic matter within the first 24 days (Opsahl and Benner 1993). By the end of the study, decomposition had resulted in a 76% loss of the organic matter. Neutral sugars were the most abundant in the starting material and also the most rapidly lost. In contrast, the most persistent sugars in the senescent blades were xylose and glucose, reflecting the more stable polymers of cellulose and xylan from which the sugars are derived. Near St. Croix, at depths between 14 and 32 m, the decomposition rates of a more delicate seagrass, H. decipiens, differed between litter bags that were buried in the sediment and those tethered at the surface of the sediment (Kenworthy et al. 1989). After 6.5 days, buried leaves lost 56% of the original ash-free dry weight compared to only 28% lost by those left on the surface. These findings were similar to two 7-day studies made in two consecutive years that included photosynthetic measurements in the same area and using the same species (Josselyn et al. 1986), this earlier

Table 4.6 δ^{34} S values for sulfur sources and plants, Redfish Bay, Texas, November 1980–February 1981 (from Fry, 1981).

Sources	δ ³⁴ S
Seawater Sulfate	+19.7 to +20.0
Sediment Sulfide	-23.4
Porewater Sulfate	+15.0 to +17.0
Rain Sulfate	$+6.3^{a}$

	Leaves		R	oots
	After			After
	Total	Washing	Total	Washing
Algae, Submerged Plants				
Digenia simplex	+18.7	+17.2		
Gracilaria verrucosa	+18.6			
Gracilaria debilis	+18.9	+17.5		
Thalassia epiphytes ^b	+17.4	+15.2		
Seagrasses, Submerged Plants				
Thalassia testudinum				
Sample 1 (November 29) ^c	+15.2		-12.3	-15.9
Sample 2 (November 29)	$+12.9^{d}$			
Sample 3 (January 3)		6.9d		-17.3
Halodule wrightii	+10.4		-9.3	
Syringodium filiforme°	+11.5		-3.3	
Halophila engelmannii	+11.2		+11.5	
Intertidal, Emergent Plants				
Spartina alterniflora	+3.5		-9.4	
Avicennia germinans ^e (as A. nitida)	-0.2		-3.2	

^{*}From Jensen and Nakai (1961).

study reporting greater than 50% decomposition of all buried seagrass material.

Seagrass epiphytes appear to be a more significant source of direct food for many invertebrates than are seagrass detritus or living tissue. Virnstein (1987) has proposed the following sequence for a south Florida seagrass food web: (1) the important primary producers are epiphytic algae, which have rapid turnover rates; (2) epiphytic algae are preferentially grazed by most species of small invertebrates associated with seagrass blades; (3) small invertebrates are preyed upon by decapod crustaceans or small fishes that include resident adults and juvenile seasonal visitors; and (4) decapods and small fishes are preyed on by larger, mostly nonresident fishes. Of the 154 grazers listed by McRoy and Helfferich (1980) that directly feed on seagrass blades, the most important invertebrates are echinoderms, molluscs, and crustaceans. For

example, seagrass blades constitute up to 50% of the total diet for crustaceans (Klumpp et al. 1989). Crabs, isopods, and sea urchins are the principal direct grazers on seagrasses, whereas other invertebrates consume seagrass detritus and various algal epiphytes. Fry and Parker (1979) and Fry (1984), using stable carbon isotopes (δ^{13} C), showed that microalgae on the sediment and seagrass blades are a major source of carbon for grazers. In some seagrass meadows, micro- and macroalgae predominate in food webs (Virnstein 1987). In a review of stable-isotope studies, Fry et al. (1987) concluded that algae are nutritionally more important for consumers than are seagrasses but that δ¹³C data alone can be misleading because benthic algae have values similar to those of seagrasses. Thus, they recommended using other elements with stable isotopes, including nitrogen and sulfur (Tables 4.5 and 4.6).

^bEpiphytic community collected after freeze drying. Contained some small animals and mud.

GePlants marked with the same letter were growing intertwined in the same sediments.

dEpiphyte-free seagrass.

RELATIONSHIPS with OTHER COMMUNITIES

Because seagrass beds are mostly found in areas of low wave energy, they often occur next to tidal-flat, salt-marsh, and mangrove communities along Florida's Gulf coast and are influenced by these intertidal communities. Salt marshes are estimated to cover about 170,000 ha (419,900 acres) of Florida's coasts; about half of the marsh area extends from Tampa Bay north to the Alabama border on the Gulf coast (Montague and Wiegert 1990). The estimated salt-marsh area is substantially less than the estimate of McNulty et al. (1972) of 214,000 ha (528,580 acres) for the Gulf coast of Florida. The largest salt marshes on the Gulf coast of Florida are in the Big Bend area, from Aripeka to Apalachicola Bay, a coast with low wave energy and extensive seagrass beds. About 60% of northwest Florida salt marshes are monotypic stands of Juncus roemerianus Scheele (Needle Rush), which often extend down to 0.2 to 0.5 m above MLW and thus are adjacent to near-shore seagrass beds. Above-ground primary production ranges from 250 to 950 g dry wt m⁻² yr⁻¹ for J. roemerianus and 130 to 700 g dry wt m⁻² yr⁻¹ for Spartina alterniflora Loisel. (Smooth Cordgrass). (Montague and Wiegert 1990), the difference probably due to the higher density of *J. roemeri*anus culms. The large biomass of salt-marsh plants and low direct consumption by herbivores (10%) means that most of the biomass decomposes and becomes part of the salt-marsh detritus (Montague et al. 1987). Apparently little of the detritus is exported to near-shore seagrass beds because of the relatively high elevation of Florida's salt marshes and often the presence of a berm that retains the detritus within the salt marsh (Montague et al. 1987).

In contrast to the low level of detrital export, a diverse fauna is shared between salt marshes and adjacent seagrass beds, for example, mullet, spot, blue crabs, oysters, and penaeid shrimps (Montague and Wiegert 1990). In addition, tarpon (Megalops atlanticus), snook (Centropomus undecimalis), red drum (Sciaenops ocellatus), seatrout (Cynoscion spp.), and kingfish (Menticirrhus spp.) move from near-shore seagrass beds into saltmarsh tidal streams (Lewis et al. 1985b).

The three Caribbean mangrove species, Rhizophora mangle L. (Red Mangrove), Avicennia germinans (L.) L. (Black Mangrove), and Laguncularia racemosa (L.) C. F. Gaertn. (White Mangrove) are estimated to cover 189,725 ha (468,620 acres) along Florida's coasts (Lewis et al. 1985b). About 90% of mangrove forests (or mangals) occur in the most southern four counties of Florida (Lee: 14,275 ha, Collier: 29,126 ha, Monroe: 94,810 ha; Dade: 32,931 ha), a region of low wave energy and consequently abundant seagrass beds. Because of the lack of severe freezing temperatures since 1989, mangroves extend intermittently northward of Tampa Bay to the western panhandle on the Gulf coast and north to the Tomoka River on the Atlantic coast (Odum and McIvor 1990; D. Crewz pers. obs.). Export of mangrove-leaf detritus is highest in riverine (1.2-2.7 g carbon m⁻² d⁻¹) and fringing forests (0.5-0.7 g carbon m⁻² d⁻¹), and these contribute to detrital-based food webs in near-shore seagrass beds (Odum and McIvor 1990). Stable carbon isotope levels (δ^{13} C) of seagrass blades and mollusk shells taken from seagrass beds near south Florida mangrove forests had significantly lower mean values (-12.8 and -2.3 ppt, respectively) than did blades and shells far from the coastal forests (-8.3 and +0.6). The lower δ¹³C values near mangrove forests indicate release of carbon dioxide by mangrove-detritus mineralization, conversion to bicarbonate, and subsequent uptake of bicarbonate by the adjacent seagrass beds (Lin et al. 1991).

The habitat values of mangals, adjacent seagrass beds, and nonvegetated open water were compared for densities of fish, shrimp, and crabs in Rookery Bay near Naples (Sheridan 1992). Fish densities were highest in nonvegetated areas (74% of the total caught), while shrimp and crab densities were highest in seagrass beds (74% and 47% of the total capture respectively). However, flooded R. mangle forests had resident and transient fish and crab numbers equal to those of other areas on some occasions. The data suggest that seagrass beds are a principal habitat for shrimp and crabs, whereas mangrove prop roots and pneumatophores may serve an ancillary role for a variety of fish and invertebrates, contrary to the conclusion drawn in earlier studies (e.g., Thayer et al. 1978, 1987).

South Florida's shallow-water coral reefs are unique in North America. Florida's reefs are simi-

lar in species composition and physiographic features to those in the Caribbean Sea and Bahama Islands (Dawes 1998a). Occasionally, hard corals occur as far north as Jupiter Island on the Atlantic coast (27°N lat.). Coral reefs form extensive threedimensional structures south and west of Cape Florida and along the Florida Keys archipelago (Jaap and Hallock 1990). Seagrass beds cover 80% of the sea bottom between Cape Sable and throughout Florida Bay, along the Florida Keys, extending westward to the Dry Tortugas, and north on the Atlantic coast into Biscayne Bay, thus co-occurring with the Florida reef tract (Jaap and Hallock 1990). Aerial photography of Biscayne National Park showed that moderately dense to highly dense seagrass beds covered 25,445 ha (62,849 acres) of the 55,000 ha (135,850 acres) in the park (Lewis et al. 2002) and were closely associated with coral patch reefs.

The relationship of seagrass beds to patch and tract reefs is considered to be in danger due to nutrient-rich runoff from canals in Biscayne Bay (Lewis et al. 2002) and from septic tanks in the Florida Keys (Jaap and Hallock 1990). A key feature of coral-reef dependence on adjacent seagrass beds is their sensitivity to nutrient enrichment, which enhances macroalgae overgrowth on the coral structures (Dawes 1998a). Thus, nutrient removal from the water column by adjacent seagrass beds may play a major role in protecting

coral reefs. In addition, the ecological role of seagrasses as habitat and nursery in the life cycle of the spiny lobster (*Panulirus argus*), whose juveniles settle in algae associated with seagrass beds, has been noted (Marx and Herrnkind 1985).

Seagrass detritus and vegetative parts are exported into deep-water habitats, especially on the Atlantic coast. Deep water (>200 m) is farther offshore on the Gulf coast of Florida than along the Atlantic, but seagrass beds are more developed in the Gulf. The larger-stature seagrass species extend to about 9 m (Dawes 1974; Iverson and Bittaker 1986). In contrast, the smaller-stature Halophila engelmannii grows in 1-5 m in Tampa Bay (Dawes 1967), in 1.4 m off Tarpon Springs (Phillips 1960b), and to 73 m off the Gulf coast (Dawes and van Breedveld 1969; Dawes and Lawrence 1990) and the Dry Tortugas (Taylor 1928). The more diminutive H. decipiens appears to be limited to deep water on the Gulf coast and occurs in 20 to 90 m (Dawes and van Breedveld 1969; Zieman 1982; Dawes and Lawrence 1990; Fonseca et al. 2001; see also Chapter 4). In the summer and early fall on the Gulf coast, H. decipiens often forms extensive beds in 20-30 m of water (Dawes and Lawrence 1990). However, no information currently exists regarding the export or role of detritus from inshore seagrass beds to the 200-m deep continental shelf west of Florida (Florida Institute of Oceanography 1994).



- Seagrass regression and die-off observed in Florida Bay in the late 1980s resulted from a combination of natural and human causes, including overproduction of seagrasses that resulted in dense shoots; anoxia that increased sediment sulfide levels; nutrient inputs that caused algal blooms, reducing light levels; lack of freshwater inputs that resulted in hypersalinity; and infection by Labyrinthula, a pathogen of Thalassia testudinum.
- The relationship between *Labyrinthula* infection and die-off of *T. testudinum* is not clear, as the pathogen has been found in a number of Florida Gulf-coast estuaries in seagrass beds that have not experienced die-off.
- Storms appear to have little direct impact on most seagrass communities, except for erosion and deposition of sediment in some areas, as found following Hurricane Andrew.
- → The 1997–1998 El Niño event caused a general die-back of seagrass communities because of low salinities and increased nutrient loads and turbidity from runoff.
- → The importance of long-shore sand bars is not adequately understood, but they may protect seagrass beds from waves and erosion in some locations.
- → Mechanical damage to seagrass beds continues, with propeller scarring being common in shallow-water communities.
- → The effects of small-scale seagrass-bed damage on local animal populations is not adequately understood; for example, moderately scarred beds show little change in shrimp and pinfish densities.
- → Nutrient pollution continues to be a serious problem, with 8 of the 15 Florida Gulf-coast estuaries regarded as being eutrophic.
- → Nutrient enrichment from natural or human sources results in increased phytoplankton density and resultant chlorophyll levels, reduction in light penetration, increased seagrass loss, increased epiphytism, increased sediment hypoxia, and shifts in seagrass species dominance.
- → The effects of sediment contaminants on seagrass communities are not clear. The highest contaminant levels of metals and pesticides are associated with fine muds, limited PAR, and low salinities.

any studies, including those by resource management programs of the National Estuary Programs for Tampa Bay, Sarasota Bay, and Charlotte Harbor and by the National Estuarine Research Reserves at Apalachicola and Rookery bays, have emphasized the importance of Florida's Gulf-coast seagrass communities. Understanding of natural- and human-induced seagrass disturbances has increased substantially, as shown in a review by Durako (1988), Short and Wyllie-Echeverria (1996) and Duarte (2002). Since 1989, many publications have examined the condition of seagrass communities on Florida's Gulf coast. These include studies of the north-central Gulf coast (Mattson 1995, 2000), Tampa Bay (Lewis et al. 1991, 1998; Johansson and Ries 1997; Kurz et al. 1999; Pribble et al. 1999; Greening 2002a; Johansson 2002a), and Sarasota Bay, Lemon Bay, and Charlotte Harbor (Kurz et al. 1999; Staugler and Ott 2000; Tomasko et al. 2001). Declines in seagrass coverage in various estuaries on the Gulf coast (including Florida Bay) have often been attributed to interactions between natural and anthropogenic factors.

In Florida Bay, *T. testudinum* die-off was noted in late 1987 by Robblee *et al.* (1991) and has been studied by many others since then (Carlson *et al.* 1994; Durako 1994; Durako and Kuss 1994; Thayer *et al.* 1994; Carlson and Yarbro 2001; Fourqurean and Robblee 1999; Hall *et al.* 1999; Zieman *et al.* 1999). Often, natural processes can result in increased damage to seagrass beds if they had been previously stressed by anthropogenic impacts, as appears to have happened regarding the seagrass die-off in Florida Bay. Despite the acknowledged interactions of natural and anthropogenic influences in these and many other systems, however, these categories are addressed separately below.

NATURAL PROCESSES

Natural processes have biological and nonbiological sources and include such influences as storms, sea urchin population increases, phytoplankton blooms, and macroalgae overgrowth. Such sources can cause direct or indirect damage that include shading, overgrazing, or disease.

Although moderate grazing is common and may even affect seagrass communities positively, overgrazing by urchins, and to a lesser extent by manatees or sea turtles, can result in complete denudation in some areas. For example, overgrazing by *Lytechinus variegatus* (purple sea urchin) resulted in severe seagrass denudation near the Pepperfish Keys (Camp *et al.* 1973) and in outer Florida Bay (Maciá and Lirman 1999; Rose *et al.* 1999).

As summarized in Chapter 4, epiphytes on seagrass blades have both positive and negative effects, with most damage being linked to shading and reductions in photosynthesis. In Tampa Bay, shading by epiphytes of *T. testudinum* blades is an important controlling factor in its depth distribution (Dixon and Leverone 1997).

High levels of drift algal biomass suppress seagrass-bed expansion and, similar to seagrass epiphytism, are usually linked to nutrient enrichment. In Hillsborough Bay, Avery (2000) reported in a 10-year study (1986–1994) that the 18 species of dominant drift algae (see Table 4.6 and Chapter 4) declined from an average biomass of 930 to 80 g dry wt m⁻² and continued to decline through 1997. In contrast, due to reductions of water-column nutrients in Hillsborough Bay, *H. wrightii* beds expanded from 0.2 ha to 40 ha (Avery 2000). On the positive side, drift and epiphytic algae probably serve as nutrient sinks and as food for a variety of invertebrates and fish (Kharlamenko *et al.* 2001; Moncreiff and Sullivan 2001).

The decline in the past 20 years of seagrass communities in Florida Bay is a conspicuous example of the effects of natural and possibly human influences on seagrass populations on the Gulf coast. In 1987, vast areas of *T. testudinum* began dying rapidly. Short-shoot densities dropped by 22% and standing crop by 28% at 107 sampling stations (Figure 5.1) between 1984 and 1994 (Hall *et al.* 1999). Losses were highest in western Florida Bay in areas with high standing crops. Similar or greater declines in standing crops were evident for *H. wrightii* (92%) and *S. filiforme* (93%) between 1984 and 1994. Hall *et al.* (1999) suggested that the most likely cause was chronic light reduction due to increased water turbidity.

A pathogen, *Labyrinthula*, was also suggested as being involved in the die-off of *T. testudinum* in Florida Bay (Porter and Muehlstein 1989; Robblee *et al.* 1991). It is similar to one that was implicated in the wasting disease of *Z. marina* over the past 50 years in the Atlantic (Burdick *et al.* 1993). Porter




Fig 5.1 Standing crop (±SD) of Thalassia testudinum in six ecological regions of Florida Bay 1984 and 1994 (Hall et al. 1999). MA = Mainland, NE = Northeast, EC = East Central, IN = Interior, AT = Atlantic, GU = Gulf.

and Muehlstein (1989) reported the presence of an undescribed species of Labvrinthula in blackened, necrotic lesions on the leaves of T. testudinum collected in seagrass die-off areas of Florida Bay. Durako and Kuss (1994) found that photosynthesis in necrotic lesions on T. testudium blades was impaired. They suggested that this may have reduced oxygen available for transport via the blade lacunae to belowground organs, resulting in increased sediment hypoxia and leading to sulfide toxicity. The pathogen has also been found to be infecting from 0% to 100% of T. testudinum blades in at least 10 sites on the Gulf coast (Figure 5.2) from the Chandeleur Islands (LA) to Florida Bay (Blakesley et al. 2002). The authors found that 60% to 90% of the sites sampled in Tampa Bay had infected short shoots of T. testudinum.

Thayer et al. (1994) ascribed the die-off in Florida Bay to "as yet unknown environmental stresses." Studies now suggest that the die-off arises from a combination of human-derived and natural factors that may collectively stress *T. testudinum* and thereby enhance infectivity by the pathogen (Blakesley et al. 2002). The factors involved include hypersalinity (Fourqurean and Robblee 1999); persistent microalgal and cyanobacterial blooms and resuspended sediments that reduced illumination (Hall et al. 1999); anoxia that increased levels of sediment sulfide (Carlson et al. 1994, 2002; Carlson and Yarbro 2001); and enhanced growth that resulted in high biomass (Zieman et al. 1999).

Hypersaline areas of Florida Bay probably resulted in part from a lack of flushing in the absence of periodic hurricanes. This condition may also be exacerbated by a reduction in freshwater flow from the Everglades because of surface-water diversion. Nevertheless, data covering 1989–1995 suggest that seagrasses were once again growing rapidly in Florida Bay and that the loss of *T. testudinum* appeared to be slowing (Zieman *et al.* 1999).

Little information is available regarding the effects of storms on Florida's Gulf-coast seagrass communities, except for that following Hurricane Andrew, which swept across southern Florida on August 24, 1992, and exited into Florida Bay at Lostmans River (Ogden 1992; Dawes et al. 1995). Although coastal mangrove forests were destroyed, seagrass beds in Lostmans Bay showed no signs of alteration or depletion of the meiofauna when sampled two and nine months after the storm (Dawes et al. 1995). To be sure, during hurricanes large volumes of water fall simultaneously upon broad landscapes, leaching large amounts of nutrients and other compounds from soils. Together with the lowering of salinities, this short-term pulse of pollutants may alter community dynamics in seagrass beds.

Also, excessive rainfall over an extended period can damage seagrass communities. An *El Niño* event between December 1997 and March 1998 resulted in over 52 cm (approximately 20 inches) of rain, which triggered widespread, persistent phytoplankton blooms, decreasing the light available to seagrasses, along the Gulf coast of Florida (Carlson *et al.* 2003). In 1997, 1998, and 1999, sampling near the Homosassa and Anclote rivers and in Tampa Bay and Charlotte Harbor found that short-shoot densities, blade widths and numbers, leaf-area indices, rhizome-apex densities, and stored carbohydrates of *T. testudinum* declined (Carlson *et al.* 2003) in response to the 1997–1998 *El Niño*.

Long-shore sand bars often occur seaward of seagrass communities and have been suggested to play a role in protecting seagrasses from strong waves and currents (Lewis 2002). Thus, reductions in seagrass coverage in some areas of Tampa Bay may reflect historical loss of these bars, similar to those in North Carolina (Fonseca and Bell 1998). In this regard to this idea, Fonseca *et al.* (2002) employed a wave-exposure model (Relative Expo-




Figure 5.2 Mean percentage of Thalassia testudinum shortshoots infected with a pathogenic species of Labyrinthula at ten sites along the Gulf coast. Numbers over bars are sample sizes. NE = not examined. (modified from Blakesley et al. 2002).

sure Index—REI) to evaluate the influence of longshore sandbars on seagrass coverage in Tampa Bay. Their results hindcasted that erosion or loss of long-shore sand bars on the southeastern shore of the bay, where wave exposure is relatively high, had caused the loss of seagrass originally located between the shore and the historical sand bars. They recommended that no restoration efforts be conducted in areas of strong wave exposure without including engineering to reduce REIs.

ANTHROPOGENIC EFFECTS

Livingston (1987) summarized anthropogenic

(human-derived) effects on seagrass meadows (Table also presented in Zieman and Zieman 1989), including direct physical damage and pollution, with an emphasis on nutrient enrichment. Since that review, various studies have described anthropogenic effects and also include estimates of seagrass losses (see Chapter 4). Nutrification and sediment loading from maintenance dredging of shipping channels and vessel-generated resuspension are cited as important causes of seagrass declines (Schoellhamer 1991; Tomasko and Lapointe 1991; Culter and Leverone 1993; Quammen and Onuf 1993; Lapointe et al. 1994). Dredging, although now generally restricted, continues in all estuaries on the Gulf coast to maintain channels and create new harbors. Certainly, such activities cause temporary resuspension of contaminated sediments and contribute to reduction of water transparency (Godcharles 1971; Schoellhamer 1991; Ailstock et al. 2002).

Mechanical damage is also done to seagrass beds by erosive effects following boat (and ship) groundings, by propeller wash, and by ship and boat wakes in shallow waters. Kenworthy et al. (1988a) found that boat wakes substantially increased bottom shear stress along edges of shallow seagrass beds. Even though nets used for bait-shrimp trawling in Tampa Bay do not appear to cause much direct damage to seagrass beds (Meyer et al. 1999) a large diversity of animals are killed as by-catch, plus trawling may suspend sediments which decreases water clarity.

Propeller scarring is another type of physical damage to seagrass beds (Figures 5.3, 5.4) that continues to increase in all areas of coastal Florida (Sargent et al. 1995). Many shallow flats and mud banks are now severely eroded due to constant scarring, ship groundings, chronic wave action from boats, and water-current scouring (Kruer 1994). Culter and Leverone (1993) stated that prop scarring was visible in almost all seagrass beds in Sarasota Bay. Sargent et al. (1995) determined that 6.2% (over 70,000 ha or 183,000 acres) of Florida's 1.1 million ha (2.7 million acres) of seagrass beds have been scarred by boat propellers (prop-dredging, prop cuts) and by similar causes, principally in coastal waters less than 2 m deep. They concluded that scarring in seagrass beds has become acute because of increasing human population densities, increasing popularities of boating, fish-



Figure 5.3 Propeller scarring from recreational boating in a Thalassia testudinum bed near Panama City, Florida.

ing, diving, and other water sports, and increasing tourism. Scarring level was divided among three categories: 1) light scarring, in which less than 5% of the seagrasses in a particular map polygon were scarred; 2) moderate scarring, in which 5% to 20% of the seagrasses within the polygon were scarred; and 3) severe scarring, in which more than 20% of the seagrasses in the polygon were scarred (Sargent et al. 1995). The highest level of scarring was found in Monroe and Citrus counties, which also contained the highest amount of seagrass coverage (688,259 ha; 1.6 million acres). Of this total, they calculated that 23,332 ha (57,630 acres) were scarred with 17.3% of the seagrasses being in Monroe County and 15.8% in Citrus County.

The influences of prop scars on seagrass community production have been studied in Tampa Bay and Charlotte Harbor (Durako et al. 1992; Clark 1995; Dawes et al. 1997; Bell et al. 2002), Sarasota Bay (Folit and Morris 1992), and the Florida Keys (Matthews et al. 1991). Fragmentation of seagrass beds in Tampa Bay due to propeller cuts did not appear to have any consistent effects on some animal populations over a oneyear period, as long as seagrass patch sizes were greater than 1 m² (Bell et al. 2002). The numbers of pinfish (L. rhomboides), pipefish (Syngnathus scovelli), and eight species of epibenthic shrimp were similar in moderately scarred (6% to 31% loss of the beds) and nonscarred seagrass beds in Tampa Bay. The results of these studies suggest that propeller scars that fragment seagrass beds may enhance certain faunal development caused by edge effects along the cuts as long as they are not too severe. Nevertheless, a recent study of scarring in a T. testudinum bed in Puerto Rico revealed a negative effect of scarring on crabs and molluscs

up to 5 m from the scar. Also, shrimp species within the scar differed from those in the non-scarred seagrasses. Fish populations did not show an effect from the scarring. Further studies are clearly needed to define the effects of moderate scarring compared to those of severe scarring on seagrass productivity.

More recent studies have confirmed previous observations by Jones (1968), Godcharles (1971), and Zieman (1976) for the long time period required for scars to heal (Eleuterius 1987; Durako et al. 1992; Dawes et al. 1997). When seagrass rhizomes and roots are completely removed by scarring, sediment is destabilized and resuspension occurs, thereby lowering water transparency and retarding seagrass regrowth into the scar. Regrowth of T. testudinum into prop scars in Tampa Bay was estimated to require an average of 3.5 to 4.1 years for existing propeller scars and up to 7.6 years in newly made ones (Dawes et al. 1997). Shorter recovery periods in existing scars probably reflect their older ages and shallower depths. Based on studies at Weedon Island (Durako et al. 1992) and Cockroach Bay (Dawes et al. 1997) in Tampa Bay, the authors recommended that shallow-water seagrass beds be protected from propeller scaring. Criteria for the protection of seagrass beds have been proposed and adopted for Tampa Bay (Clark 1995; Stowers et al. 2002).

Another source of physical impacts that degrade seagrass communities is docks, which shade seagrass beds, as shown in Perdido Bay (Shafer 1999), Charlotte Harbor (Loflin 1995), and the Indian River Lagoon (Beal and Schmit 2000). Smith and Mezich (1999) surveyed 200 out of the 3,592 permit-exempt single- and multi-family docks that were deemed to have the potential to




Figure 5.4 Propeller scarring from commercial fishing in a seagrass bed in Pine Island Sound, Florida.

damage seagrass beds in Palm Beach County. They found that 16% of the docks were larger than regulations allowed, owing to additions, and that approximately 1,491 single-family docks in the county had affected seagrass negatively. They estimated that, in total, docks eliminated 20.4 ha (50.4 acres) of seagrass beds, which is equal to 2.1% of the seagrass extents in Palm Beach County. Their recommendations were that all docks in Florida be considered within the jurisdiction of FDEP permitting programs. Currently, environmental resource permits are not required within designated Aquatic Preserves if the docks are less than 46.5 m² (500 ft²) or if they are less than 92.9 m² (1,000 ft.2) when outside Aquatic Preserves. Smith (1998) concluded that strong enforcement of the regulations protecting sovereign submerged lands (253.04-05, 380.05, F.S. and Chapter 18-14, F.A.C.) would act as a powerful deterrent against boat and dock damage to seagrass systems.

Direct physical damage of seagrasses can result from changes in freshwater inputs. The importance to watershed management and reduction of freshwater removal from tributaries flowing into Gulf-coast estuaries has been reviewed for Tampa Bay (Zarbock 1991) and for Charlotte Harbor (Kurz et al. 1999). Estevez (2000) found that changes in the amount, timing, and location of freshwater inflow are primary stressors to estuarine and oceanic seagrass communities, with salinity changes often being a first-order stressor (Montague and Ley 1993). The present estimated daily freshwater inflows into Tampa Bay (Zarbock 1991) range from 152.0 to 214.6 m³ s⁻¹ (1,792 to 2,530 ft³ s⁻¹) compared to historical estimates of 193.4 m³ s⁻¹ (2,280 ft³ s⁻¹). Although timing and duration of riverine inflow, along with nutrient

loading and changes in water color, clearly affect seagrass communities in Charlotte Harbor, the relationships between these factors is more complex in Tampa Bay and Sarasota Bay (Kurz et al. 1999). Similarly, diversion of fresh water before it enters the Everglades, coupled with a decrease in hurricane-induced flushing (Thayer et al. 1994) and rising salinities and sediment anoxia (Zieman et al. 1999), has been proposed as influencing factors in the precipitous decline of Florida Bay seagrasses.

Short and Neckles (1999) hypothesized that global warming will probably influence seagrass distribution by causing increases in sea level, in storm frequency and intensity, in disease from higher water temperatures, and in turbidity from eutrophication. The authors also predicted an increase in ultraviolet radiation that will damage seagrasses, as has been shown for macroalgae (Larkum and Wood 1993) and terrestrial plants (Caldwell et al. 1989). In addition, the shallow coastal and estuarine waters and low elevations on the Gulf coast of Florida will certainly be affected by any eustatic sea-level rise. A potential positive effect on seagrass distribution may occur from an increase in dissolved inorganic carbon (Ci) composition in seawater, as seagrasses are presently CO2 limited and have an affinity for Ci (Beer and Koch 1996).

Nutrient enrichment, probably the most common human effect in Gulf-coast estuaries, can result in a variety of changes in seagrass communities. A number of these changes have been presented in Chapter 4. These include an increase in water-column chlorophyll levels (Johansson 1991; Janicki *et al.* 1999; Tomasko *et al.* 1996, 2001; Morrison *et al.* 1997), in algal epiphytism and shading

of seagrass blades (Tomasko and Lapointe 1991; Tomasko *et al.* 1996; Dixon 2002), in shifts in dominant seagrass species (Duarte 1995; Fourqurean *et al.* 1995), and in sediment hypoxia (Carlson *et al.* 2002).

A review of U.S. estuaries (Bricker et al. 1999) cited 15 estuaries on the Gulf coast of Florida that had various levels of eutrophic conditions. Eight of these estuaries (Florida Bay, the southern Ten Thousand Islands, Caloosahatchee River, Charlotte Harbor, Sarasota Bay, Tampa Bay, Choctawhatchee Bay, and Perdido Bay) had high levels of eutrophic symptoms. More detailed data are available regarding nutrient loadings and damage to seagrasses in Tampa Bay (Treat and Clark 1991; Zarbock et al. 1994, 1996; Treat 1997; Lewis et al. 1998; Pribble et al. 1999; Greening 2002b), Sarasota Bay (Haddad 1989; Tomasko et al. 1992, 1996), and Lemon Bay (Tomasko et al. 2001). Tampa Bay has been the subject of several reviews (Treat and Clark 1991; Treat 1997; Greening 2002a) that include data regarding point- and nonpoint-source pollution, domestic and industrial effluents, accumulation of toxic substances in the sediment, and potential brine damage from desalination facilities.

Many studies worldwide support the contention that seagrass distribution, survival, and growth are principally determined by water clarity (Batiuk et al. 1992; Dennison et al. 1993; Duarte 1991; Kenworthy and Haunert 1991; Kenworthy 1993; Koch and Beer 1996; Morris and Tomasko 1993; and Tomasko et al. 2001; also individual papers contained in the publications of Bortone 2000 and Greening 2002a). However, other factors such as waves and water currents can affect seagrass distribution as well (Koch 2001; see Chapter 2). Water clarity is influenced by nutrient inputs, which stimulate the growth of phytoplankton (particularly by nitrogen), and by total dissolved and suspended solids. An overabundance of nutrients leads to eutrophication that supports increased levels of phytoplankton, drift algae biomass, and coverage by epiphytes on seagrass blades, all of which result in shading of seagrass leaves (Sand-Jensen 1977, 1990; Cambridge et al. 1986). In all cases a decrease in water clarity over seagrass beds was correlated with a decline in seagrass standing stock (Orth and Moore 1983; Short et al. 1996; Short and Wyllie-Echeverria 1996; Fitzpatrick and Kirkman 1995; Lewis et al. 1985a).

The influence of possible sediment contaminants on seagrass beds is not understood. The concentration of metals, nutrients, petroleum, chlorinated hydrocarbons, and radionuclides correlate with the smaller particle-size sediments (muds; <63µm in diam.) in Tampa Bay (Brooks and Doyle 1991) and with sediment-quality factors (Zarbock et al. 1997). Thus, concentrations of all contaminants are highest in west-central Hillsborough Bay and Old Tampa Bay (Brooks and Doyle 1991) for total organic matter (to 16%), total hydrocarbons (>40 µg g⁻¹), total organic nitrogen (0.1%), and total phosphorus (0.4%). Using the triad method of weighting sediment chemistry, toxicity, and benthic samples, Zarbock et al. (1997) determined that upper and middle Hillsborough Bay, parts of Old Tampa Bay, Boca Ciega Bay, and western Middle Tampa Bay had the most contaminated sediments. In a review of sediment quality in Tampa Bay, Grabe (1999) found that approximately 1% of its sediments were subnominal and had a high probability of being toxic. However, at the levels known for Tampa Bay, evidence that the heavy metals and pesticides directly affect seagrasses does not exist.

In conclusion, it is apparent that a mix of natural and human-induced events has negatively influenced the seagrass meadows on the Gulf coast of Florida. Because of limited historical data, determining the extent of seagrass decline is often difficult, although detailed historical analyses are possible in some areas, such as for Tampa Bay (Greening 2002a). Anthropogenic influences, particularly those that result in a decline in water quality, are considered the principal causes of seagrass losses throughout the world (Short and Wyllie-Echeverria 1996) and in the estuaries of Florida's Gulf coast (Lewis et al. 1985a). The loss of seagrass communities continues worldwide (Duarte 2002) as well as on Florida's Gulf coast (see Chapter 2). However, restoration management can result in the expansion and enhancement of seagrass meadows, as seen in Tampa Bay and Sarasota Bay (Greening 2002b). Future studies are needed to distinguish secondary and primary stressors that result in seagrass declines, such as the modification of watershed-level stormwater runoff and direct nutrient input.

SAXONO FEORIDA SEAGRASSES

APPENDIX

Six seagrass species, known throughbut the Caribbean Sea and Gulf of Mexico, occur on the Gulf coast of Florida. A seventh species, Halophila johnsonii, is known only from the Atlantic coast of Florida. Worldwide, sixty species of seagrasses are recognized. They are currently placed by Tomlinson (1982) in 12 genera and four families in a single class of monocots, the Liliopsidt (Kuo and McComb 1989) and one subclass, Helphiae (Tomlinson 1982). The line usion of the presentation as a seagrass (Dawes 1998). Littler and Littler 2000) brings the monocot of species to 61 and families to 5. The family and genus descriptions medified from Dawson (1966), Tomlinson (1982), and Kurana men Hartog (2001). A key based on vegetative characters is followed by the set descriptions modified from den Hartog (1970), Littler and Littler (2000), and Kurana and den Hartog (2001).

Charles Seaborn photo

ARTIFICIAL KEY to SPECIES, BASED on VEGETATIVE FEATURES

HYDROCHARITACEAE Jussieu

Family of 15 genera, diverse and cosmopolitan, of which 3 are seagrasses (*Enhalus*, *Halophila*, *Thalassia*). Plants aquatic herbs, leaves either large, linear to strap-shaped and with sheathing base (*Thalassia*) or small, usually nonlinear, with distinct petiole and without sheathing base (*Halophila*). Rhizomes with scale-like leaves, and branching via apical bifurcation. Flowers unisexual and anthers either sessile or on long slender filaments. Female flowers usually with long hypanthium and inferior ovary. Fruits fleshy and often indehiscent.

THALASSIA Banks and Solander ex König. 1805. Ann. Bot. (König et Sims) 2: 96 (Gr. thalassa: marine, of the sea).

Rhizomes creeping, horizontal, thick, fleshy, and indeterminate; bearing scale leaves and erect short-shoots (rhizomes) at irregular intervals that arise from rhizome meristem. Erect short shoots determinate, encased in leaf sheaths, producing foliate leaves and flowers. Roots unbranched, fleshy, arising at nodes of short shoots. Leaves strap-shaped, with sheathing bases, and growing from basal meristem. Short shoots with conspicuous leaf and inflorescence scars. Plants are dioecious. Male flowers on short stalks, in clusters of 1–3. Female flowers solitary. Fruits globose and

opening by irregular splitting, containing several large angular "seeds" (seedlings).

Thalassia testudinum Banks and Solander ex König. 1805. Ann. Bot. (König et Sims) 2: 96 (L. testudinis: a tortoise). Common name: Turtlegrass (Figure A.1.4).

Rhizomes creeping, 3–6 mm in diameter, 4–7 mm long between nodes, with scale leaves. Roots single, unbranched, arising at short-shoot nodes, producing delicate root hairs at tips. Erect short shoots with clusters of 3–7 leaves; leaves to 2 cm wide, with persistent colorless basal sheath, 9–17 parallel veins, and lacunae that are continuous throughout the plant. Plants dioecious. Flowers on short peduncles, with single whorl of white tepals. Male flowers with 3 to 12 stamens, pollen grains in mucilage, forming moniliform chains. Female flowers bearing one ovary with 6–8 carpels; style divided into 2 filiform stigmata. Fruits buoyant, 15–20 mm long, rough, fleshy, tips pointed, with one to few "seeds" (seedlings).

HALOPHILA Du Petit-Thouars. 1806. Nova Madag. 2 (L. *halo*: salt + *philus*: loving).

Rhizomes horizontal, usually on surface and bearing two scales at each node, producing a lateral shoot and one unbranched root, short erect shoots either with paired leaves on long petioles, distichously arranged along the axes, or in a pseudo-whorl at top of shoot. Leaves oval, elliptical, lanceolate or linear, with mid and marginal veins. Plants monoecious or dioecious. Flowers covered by 2 bracts; sepals small, petals absent. Male flowers stalked, with 3 stamens having sessile anthers. Female flowers sessile, with 3–6 styles. Fruits ovoid capsules, bearing many seeds.

Halophila decipiens **Ostenfeld.** 1902. Bot. Tildsskr. 24: 260 (L. *decipio*: beguiling, deceptive). Common name: Paddlegrass (Figure A.1.1).

Rhizomes thin, with transparent, hairy scales, internodes 1.0–4.5 cm long. Petioles 3–15 mm long, bearing one pair of leaves; leaves oblong to elliptical, 10–25 mm long, 3–6 mm wide, with short rigid unicellular hairs on one or both sides of blades, margins finely serrate. Plants monoecious, male and female flowers on same stalk, covered by spathe, stalks to 1.5 mm long. Male flowers with 2 anthers to 1 mm long. Female flowers with 3 styles 1.5–2.5 mm long; single ovary to 1 mm long. Fruits 2.5 mm long, 1.5 mm wide, elliptical, bearing up to 30 seeds; seeds oval, to 0.2 mm long.

Halophila engelmannii Ascherson in Neumayer. 1875. Anl. Wiss. Beobeibet Reiser. p. 368 (Named for George Engelmann [1809–1885], an American physician and German botanist who settled in St. Louis). Common name: Stargrass (Figure A.1.3).

Rhizomes indeterminate, narrow, with internodes 2–4 cm long. Determinate, erect shoots 20–40 mm tall at each node. Leaves with petioles, 2 scales at base and another pair half-way up shoot, petioles to 2 mm long, bearing 2–4 pairs of leaves in pseudo-whorls. Blades oblong to linear-oblong, 10–30 mm long, 3–6 mm wide, tips pointed and margins serrate. Plants dioecious. Male flowers with 3 imbricate tepals and 3 stamens. Female flowers with one sessile, inferior ovary, ovaries 3–4 mm long, ovoid, with 1–3 styles; styles to 30 mm long. Fruit globose, fleshy capsule, 3–4 mm in diam. with several minute subspherical seeds.

Halophila johnsonii Eiseman. 1980. In Eiseman and McMillan, Aquatic Bot. 9: 16 (Named for J. Seward Johnson, cofounder of Harbor Branch Oceanographic Institute, where N. Eiseman studied). Common Name: Johnson's Seagrass. (Figure A.1.2).

Rhizomes thin, 1 mm in diam., nodes bearing scale leaves on upper and lower surfaces, internodes 1–2 cm long. Petioles 10–20 mm long, bear-

ing pair of linear to spatulate leaves; leaves slightly asymmetrical, 5–25 mm long, 1–4 mm wide, margins entire and surfaces glabrous. Plants dioecious (?), only female plants found. Female flowers sessile, with three styles 4–6 mm long; seeds unknown. occurring only in Atlantic Florida (Sebastian Inlet to Biscayne Bay) and classified as a threatened species. May not be distinct from *H. ovalis* (R. Brown) J.D. Hooker *f.*, 1858.

CYMODOCEACEAE N. Taylor

Family of 5 morphologically distinct genera and about 20 species, all seagrasses. Plants perennial herbs. Rhizomes creeping, leafy, herbaceous or distinctly woody, scale-bearing. Leaves distichous with distinct sheath and blade; blades linear, flat or terete, leaves with numerous tannin cells. Plants dioecious. Flowers solitary or in cymose inflorescences and usually terminal. Male flowers subsessile or stalked with 2 anthers producing filiform pollen. Female flowers with 2 free carpels, each with 1 style. Fruits 1-seeded nuts, indehiscent.

HALODULE Endlicher. 1841. Gen. 1368 (L. halo: salt + dule: loving).

Rhizomes monopodially branched; each node producing one to several unbranched roots and erect short shoot with 1–4 leaves. Leaves narrow, linear, with 3 longitudinal veins and short basal sheaths. Plants are dioecious. Flowers solitary, terminal, enclosed by leaf. Male flowers stalked, with 2 anthers. Female flowers with 2 free carpels, with long undivided styles. Fruits oval to slightly flattened, with a stony pericarp.

Halodule wrightii Ascherson. 1868. Bot. Zeitung (Berlin) 26: 511 (Named for Charles Wright [1811–1885], who collected extensively in the southwest U.S. and Cuba and sent specimens to Ascherson). Common names: Shoalgrass, Cuban Shoalgrass, or Shoalweed (Figure A.2.3).

Rhizomes slender, bearing 2–4 roots and a short shoot at each node; internodes 0.75–3.5 cm long, with elliptic scale leaves 5–10 mm long. Leaf sheaths 1.5–6.0 cm long; blades 5–12 (–20) cm long, 0.50–1.5 (–2.0) mm wide, with 3 parallel veins; blade tips with 2–3 short horn-like points. Plants are dioecious. Flowers lack tepals; male flowers on peduncles 1.5–2.0 cm long, anthers slender, 3.5–5.0 mm long. Female flowers with one

oval to elliptic ovary, 1.5–2 mm wide, with 10–28 mm long style. Fruits about 1.5–2.0 mm in diam. with stony pericarp and stylar beak.

SYRINGODIUM Kützing. 1860. In Hohenacker, Alg. Marin. Sicc. 9: 426 (Gr. *Syringx*: a tube + *odium*: similar to; thus like *Syringa*, a genus of flowering plants).

Rhizomes indeterminate, with monopodial branching and scale leaves; each node bearing 1–3 branching roots and erect, unbranched, determinate short shoot. Short shoots producing 2–3 cylindrical leaves with basal sheaths. Plants dioecious. Flowers in erect cymose infloresences; male flowers stalked, with 2 anthers. Female flowers with 2 free carpels, each with short style. Fruits with stony, smooth pericarp.

Syringodium filiforme Kützing. 1860. In Hohenacker, Alg. Marin. Sicc. 9: 426 (L. *filiforme:* long and thin). Common name: Manateegrass (Figure A.2.2).

Rhizomes 2–4 mm in diameter, nodes bearing 2–4 roots with abundant root hairs, scale leaves, and erect short shoot. Short shoots with 2–3 leaves; leaves to 30 cm long, 1–2 mm in diam., cylindrical, with extensive basal sheathing, cylindrical. Plants dioecious. Male flowers on short peduncles, with 2 anthers. Female flowers sessile, with leaf sheath and 2 free carpels. Fruits oval to oblique, to 6 mm long, 3–5 mm in diam.

RUPPIACEAE Hutchins

Family of one genus and 2–3 species, all seagrasses, closely related to the Potomogetonaceae. Plants glabrous, submerged perennial herbs found in

alkaline lakes and brackish to oceanic waters. Rhizomes creeping and little differentiated from erect axes, with monopodial branching. Leaves opposite or alternate, narrowly linear, with single medial vein and short basal sheath. Plants monoecious. Flowers perfect, on spikes subtended by 2 subopposite involucral leaves with slightly inflated sheaths; two flowers per spike, each with 2 stamens and few to 4 carpels with short, stout to finely attenuate styles and peltate stigmas. Fruits indehiscent achenes, the outer layers soft, decaying and inner pericarp stony.

RUPPIA Linnaeus. 1753. Species Plantarum, p. 127 (Named for Heinrich Reinhard Ruppius [1688–1719], a German botanist who wrote the Flora of Jena).

Genus with characters of the family.

Ruppia maritima Linnaeus. 1753. Species Plantarum, p. 127 (L. *maritima*: of the ocean, marine). Common names: Ditchgrass; Widgeongrass (Figure A.2.1).

Rhizomes thin, 1–3 mm in diameter, profusely branched, with growth forms; often geniculate. Leaves produced at nodes, sheath 6–10 mm long. Reproductive axes on erect stalks 60–90 cm long. Leaves to 1 dm long, 0.3 mm wide, 1 veined, with pointed tips and appearing to arise directly from rhizome. Plants monoecious. Flowers on short (>6 cm) peduncles elongating after anthesis, becoming loosely coiled spiral. Flowers with 2 sessile anthers 2-celled, elliptical, 0.6–0.7 mm long; carpels oval, to 2 mm long, style short and stout to attenuate. Fruitlets (nutlets) 2–3 mm long, on short stipe, often curved.




Figure A.1 HYDROCHARITACEAE: 1 Halophila decipiens, Paddlegrass (x 2/3); 2 H. johnsonii, Johnson's Seagrass (x 3/4); 3 H. engelmannii, Stargrass (x 3/4); 4 Thalassia testudinum, Turtlegrass (x 1/3). Scale approximate. Smaller phenotypes are common.




Figure A.2 RUPPIACEAE: 1 Ruppia maritima, Widgeongrass (\times 1/4). CYMODOCEACEAE: 2 Syringodium filiforme; Manateegrass (\times 1/4); 3 Halodule wrightii, Shoalgrass (\times 1/3). Scale approximate. Smaller phenotypes are common.

REFERENCES

- Abel, K.M. and E.A. Drew. 1989. Carbon metabolism. Pp. 760–796 in Larkum, A.W.D., A.J. McComb, and S.A. Shepherd (eds.), Biology of Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region. Amsterdam, The Netherlands. 841 pp.
- Adams, J.B. and G.C. Bate. 1994. The ecological implications of tolerance to salinity by *Ruppia cirrhosa* (Petagna) Grande and *Zostera capensis* Setchell. *Botanica Marina* 37:449–456.
- Agami, M. and Y. Waisel. 1988. The role of fish in distribution and germination of seeds of the submerged macrophytes *Najas marina* L. and *Ruppia maritima* L. *Oecologia* 76:83–88.
- Ailstock, M.S., S.G. Hornor, C.M. Norman, and E.M. Davids. 2002. Resuspension of sediments by water-craft operated in shallow water habitats of Anne Arundel County, Maryland. *Journal of Coastal Research* (Special Issue) 37:18–32.
- Ancibar, E. 1979. Systematic anatomy of vegetative organs of the Hydrocharitaceae. *Botanical Journal of the Linnaean Society* 78:237–266.
- Andorfer, J. H. 2000. Physiological integration and translocation of carbon and nitrogen in the seagrass *Thalassia testudinum* Banks ex König. Ph.D. Dissertation, University of South Florida, Tampa, Florida.
- Andorfer, J. and C.J. Dawes. 2002. Production of rhizome meristems by the tropical seagrass *Thalassia testudinum*: the basis for slow recovery into propeller scars. *Journal of Coastal Research* (Special Issue) 37:130–142.
- Andrews, T.J. and K.M. Abel. 1977. Photosynthetic carbon metabolism in seagrasses: ¹⁴C labeling evidence for the C₃ pathway. *Plant Physiology* 63:650–656.
- Angel, R. 2002. Genetic diversity of *Halodule wrightii* using random amplified polymorphic DNA. *Aquatic Botany* 74:165–174.
- Arber, A. 1920. Water Plants: A Study of Aquatic Angiosperms. Cambridge University Press, London, England. 436 pp. plus illustrations.
- Avery, W.M. 1997. Distribution and abundance of macroalgae and seagrass in Hillsborough Bay, Florida, from 1986 to 1995. Pp.151–166 in Treat, S.F. (ed.), Proceedings: Tampa Bay Area Scientific Information Symposium 3: Applying Our Knowledge.

- Tampa Bay Regional Planning Council, Tampa, Florida. 396 pp.
- Avery, W.M. 2000. Monitoring submerged aquatic vegetation in Hillsborough Bay, Florida. Pp. 137–145 in Bortone, S.A. (ed.), Seagrasses: Monitoring, Ecology, Physiology, and Management. CRC Press, Boca Raton, Florida. 318 pp.
- Ballentine, D.L., N.E. Aponte, and J.G. Holmquist. 1994. Multi-species algal balls and potentially imprisoned fauna: an unusual benthic assemblage. *Aquatic Botany* 48:167–174.
- Barber, B.J. and P.J. Behrens. 1985. Effects of elevated temperature on seasonal *in situ* leaf productivity of *Thalassia testudinum* Banks ex König and *Syringodium filiforme* Kützing. *Aquatic Botany* 22:61–69.
- Batiuk, R.A., R.J. Orth, K.A. Moore, W.C. Dennison, J.C. Stevenson, L.W. Staver, V. Carter, N.B. Rybicki, R.E. Hickman, S. Kollar, S. Bieber, and P. Heasly. 1992. Chesapeake Bay submerged aquatic vegetation habitat requirements and restoration targets: a technical synthesis. CBP/TRS 83/92. Environmental Protection Agency and Chesapeake Bay Program, Annapolis, Maryland. 186 pp.
- Baydoun, E. A.-H. and C.T. Brett. 1985. Comparison of cell wall compositions of the rhizomes of three seagrasses. *Aquatic Botany* 23:191–196.
- Beal, J.L. and B.S. Schmit. 2000. The effects of dock height on light irradiance (PAR) and seagrass (Halodule wrightii and Syringodium filiforme) cover. Pp. 49–63 in Bortone, S.A. (ed.), Seagrasses: Monitoring, Ecology, Physiology, and Management. CRC Press, Boca Raton, Florida. 318 pp.
- Beer, S. 1996. Inorganic carbon transport in seagrasses. Pp. 43–47 in Kuo, J., R.C. Phillips, D.I. Walker, and H. Kirkman (eds.), Seagrass Biology: Proceedings of an International Workshop, Rottnest Island (2 Vols.), University of Western Australia, Nedlands, Western Australia. 663 pp.
- Beer, S., M. Björk, F. Hellbloom, and L. Axelsson. 2003. Inorganic carbon utilization in marine angiosperms. *Functional Plant Biology* 29:349–354.
- Beer, S. and E. Koch. 1996. Photosynthesis of marine macroalgae and seagrasses in globally changing CO₂ environments. *Marine Ecology Progress Series* 141:199–204.
- Beer, S., A. Eschel, and Y. Waisel. 1977. Carbon metab-

- olism in seagrasses. 1. Utilization of exogenous inorganic carbon species in photosynthesis. *Journal of Experimental Botany* 28:1180–1189.
- Beer, S., A. Shomer-Ilan, and Y. Waisel. 1980. Carbon metabolism in seagrasses. 2. Patterns of photosynthetic CO₂ incorporation. *Journal of Experimental Botany* 31:1019–1026.
- Bell, F.W. 1993. Current and projected tourist demand for saltwater recreational fisheries in Florida. SGR-111. Florida Sea Grant College Program, Gainesville, Florida. 95 pp.
- Bell, S.S., R.A. Brooks, B.D. Robbins, M.S. Fonseca, and M.O. Hall. 2001. Faunal response to fragmentation in seagrass habitats: implications for seagrass conservation. *Biological Conservation* 100:115–123.
- Bell, S.S., L.A.J. Clements, and J. Kurdziel. 1993. Production in natural and restored seagrasses: a case study of a macrobenthic polychaete. *Ecological Applications* 3:610–621.
- Bell, S.S., M.O. Hall, and M.S. Fonseca. 1994. Evaluation of faunal and floral attributes of seagrass beds in high and low energy regimes: a geographic comparison. Pp. 267–272 in Dyer, K.R. and R.J. Orth (eds.), Changes in Fluxes in Estuaries: Implications from Science to Management. ECSA22/ERF Symposium. Olsen and Olsen Publishers, Fredensborg, Denmark. 496 pp.
- Bell, S.S., M.O. Hall, and B.D. Robbins. 1995. Toward a landscape approach in seagrass beds—Using macroalgal accumulation to address questions of scale. *Oecologia* 104:163–168.
- Bell, S.S., M.O. Hall, S. Soffian, and K. Madley. 2002. Assessing the impact of boat propeller scars on fish and shrimp utilizing seagrass beds. *Ecological Appli*cations 12:206–217.
- Benedict, C.R. and J.R. Scott. 1976. Photosynthetic carbon metabolism in a marine grass. *Plant Physiology* 57:876–880.
- Benedict, C.R., W.W.L. Wong, and J.H.H. Wong. 1980. Fractionation of the stable isotopes of inorganic carbon by seagrasses. *Plant Physiology* 65:512–517.
- Björk, M., J. Uku, A. Weil, and S. Beer. 1999. Photosynthetic tolerances to desiccation of tropical intertidal seagrasses. *Marine Ecology Progress Series* 191:121–126.
- Blakesley, B.A., D.M. Berns, M.F. Merello, M.O. Hall, and J. Hyniova. 2002. The dynamics and distribution of the slime mold *Labyrinthula* sp. and its

- potential impacts on *Thalassia testudinum* populations in Florida. Pp. 199–207 in Greening, H.S. (ed.), *Proceedings, Seagrass Management: It's Not Just Nutrients!* Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Bortone, S.A. (ed.). 2000. Seagrasses: Monitoring, Ecology, Physiology, and Management. CRC Press, Boca Raton, Florida. 318 pp.
- Bottone, P.J. and R.A. Mattson. 1987. Preliminary report on transplanting of the benthic green alga Caulerpa prolifera. Pp. 63–74 in Webb, F.J. (ed.), Proceedings of the Fourteenth Annual Conference on Wetlands Restoration and Creation. Hillsborough Community College, Tampa, Florida.
- Bottone, P.J. and D.M. Savercool. 1993. Competition for space between *Caulerpa prolifera* and *Halodule wrightii* in Tampa Bay. Final Report. Peninsula Design and Engineering, Inc., Tampa, Florida.
- Bricker, S.B., C.G. Clement, D.E. Pirhalla, S.P. Orlando, and D.R.G. Farrow. 1999. National estuarine eutrophication assessment. effects of nutrient enrichment in the nation's estuaries. National Oceanic and Atmospheric Administration, National Ocean Service, Special Projects Office and the National Centers for Coastal Ocean Science, Silver Spring, Maryland.
- Bridges, K.W., J.C. Zieman, and C.P. McRoy. 1978. Seagrass literature survey. *Degraded Material Research Program Report D-78-4*. U.S. Army Corps of Engineers, Waterways Experiment Station, Vicksburg, Mississippi. 174 pp.
- Brook, I.M. 1977. Trophic relationships in a seagrass community (*Thalassia testudinum*) in Card Sound, Florida. Fish diets in relation to macrobenthic and cryptic faunal abundance. *Transactions of the American Fisheries Society* 106:219–229.
- Brook, I.M. 1978. Comparative macrofaunal abundance in turtle grass (*Thalassia testudinum*), in south Florida characterized by high blade density. *Bulletin of Marine Science* 28:212–217.
- Brooks, G.R. and L.J. Doyle. 1991. Distribution of sediments and sedimentary contaminants in Tampa Bay. Pp. 399–414 in Treat, S.F. and P.A. Clark (eds.), Proceedings, Tampa Bay Area Scientific Information Symposium 2: The Watershed. TEXT, Tampa, Florida. 528 pp.
- Brooks, R.A. and S.S. Bell. 2001. Mobile corridors in marine landscapes: enhancement of faunal exchange at seagrass/sand ecotones. *Journal of Experimental Marine Biology and Ecology* 264:67–84.

- Brown, M. M. 2001. The drift algae of seagrass beds in Sunset Beach, Cockroach Bay, and Tarpon Bay on the west coast of Florida. M.S. Thesis, University of South Florida, Tampa, Florida.
- Burdick, D.M., F.T. Short, and J. Wolf. 1993. An index to assess and monitor the progression of wasting disease in eelgrass *Zostera marina*. *Marine Ecology Progress Series* 94:83–90.
- Cain, M.L. 1994. Consequences of foraging in clonal plant species. *Ecology* 75:933–944.
- Caldwell, M.M., A.H. Teramura, and M. Tevini. 1989. The changing solar ultraviolet climate and the ecological consequences for higher plants. *TREE* 4:363–367.
- Cambridge, M.L., A.W. Chiffings, C. Brittan, L. Moore, and A.J. McComb. 1986. The loss of seagrass in Cockburn Sound, Western Australia. II. Possible causes of seagrass decline. *Aquatic Botany* 24:269–285.
- Camp, D.K., S.P. Cobb, and J.F. van Breedveld. 1973. Overgrazing of seagrasses by a regular urchin *Lytechinus variegatus*. *BioScience* 23:37–38.
- Carlson, P.R., Jr., and L.A. Yarbro. 2001. Seagrass mortality and recovery on Florida Bay mudbanks. Final Report for Cooperative Agreement CA5280-8-5006. Florida Marine Research Institute, St. Petersburg, Florida.
- Carlson, P.R., Jr., L.A. Yarbro, and T.R. Barber. 1994. Relationship of sediment sulfide to mortality of *Thalassia testudinum* in Florida Bay. *Bulletin of Marine Science* 54:733–746.
- Carlson, P., L. Yarbro, T. Barber, and D. Smith. 1999. A conceptual model of Florida Bay seagrass mortality (1987–1991): links between climate, circulation, and sediment toxicity. *Abstract*, Pp. 10–11 *in* Florida Bay and Adjacent Marine Systems Science Conference.
- Carlson, P.R., L.A. Yarbro, K. Madley, H. Arnold, M. Merello, L. Vanderbloemen, G. McRae, and M.J. Durako. 2003. Effect of El Niño on demographic, morphological, and chemical parameters in turtle-grass (*Thalassia testudinum*): an unexpected test of indicators. *Journal of Environmental Monitoring and Assessment* 81:393–408.
- Carlson, P.R., Jr., L.A. Yarbro, B.J. Peterson, A. Ketron, H. Arnold, and K.A. Madley. 2002. The influence of sediment sulfide on the structure of south Florida seagrass communities. Pp. 215–217 in Greening, H.S. (ed.), *Proceedings, Seagrass Management: It's Not Just Nutrients!* Tampa Bay Estuary Program, St.

- Petersburg, Florida. 246 pp.
- Cebrián, J., C.M. Duarte, N.S.R. Agawin, and M. Merino. 1998. Leaf growth response to simulated herbivory: a comparison among seagrass species. *Journal of Experimental Marine Biology and Ecology* 220:67–81.
- Christian, D. and Y.P. Sheng. 2002. Light attenuation by color, chlorophyll a, and tripton in Indian River Lagoon. Pp. 91–105 in Greening, H.S. (ed.), *Proceedings, Seagrass Management: It's Not Just Nutrients!* Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Clark, P.A. 1995. Evaluation and management of propeller damage to seagrass beds in Tampa Bay, Florida. *Florida Scientist* 58:193–197.
- Collado-Vides, L., G. González-González, and M.G. Morgan. 1994. A descriptive approach to the floating masses of algae of a Mexican Caribbean coastal lagoon. *Botanica Marina* 37:391–396.
- Costanza, R. 1999. The ecological, economic, and social importance of the oceans. *Ecological Economics* 31:199–213.
- Costanza, R., R. d'Arget, R. de Groot, S. Farber, M. Grasso, B. Hannon, K. Limburg, S. Naeom, R.V. O'Neill, J. Paruelo, R.G. Raskin, P. Sutton, and M. van Belt. 1997. The value of the world's ecosystem services and natural capital. *Nature* 387:253–260.
- Cox, P.A. and C.J. Humphries. 1993. Hydrophilous pollination and breeding system evolution in seagrasses: a phylogenetic approach to the evolutionary ecology of the Cymodoceaceae. *Botanical Journal of the Linnaean Society* 113:217–226.
- Cox, P.A. and P.B. Tomlinson. 1988. Pollination ecology of a seagrass, *Thalassia testudinum* (Hydrocharitaceae), in St. Croix. *American Journal of Botany* 75:958–965.
- Culter, J.K. and J.R. Leverone. 1993. Bay bottom habitat assessment. *Mote Marine Laboratory Technical Report No. 303*. Mote Marine Laboratory, Sarasota, Florida. 69 pp. plus appendixes.
- Czerny, A.B. and K.H. Dunton. 1995. The effects of *in situ* light reduction on the growth of two subtropical seagrasses, *Thalassia testudinum* and *Halodule wrightii*. Estuaries 18:418–427.
- Davis, J.L., D.L. Childers, and D.N. Kuhn. 1999. Clonal variation in a Florida Bay *Thalassia testudinum* meadow: molecular genetic assessment of population structure. *Marine Ecology Progress Series* 186:127–136.
- Dawes, C.J. 1967. Marine Algae in the Vicinity of

- *Tampa Bay, Florida.* University of South Florida, Tampa, Florida. 104 pp.
- Dawes, C.J. 1974. Marine Algae of the West Coast of Florida. University of Miami Press, Coral Gables, Florida. 201 pp.
- Dawes, C.J. 1986. Seasonal proximate constituents and caloric values in seagrasses and algae on the west coast of Florida. *Journal of Coastal Research* 2:25–32.
- Dawes, C.J. 1998a. *Marine Botany*, 2nd Ed. John Wiley and Sons, Inc., New York, New York. 480 pp.
- Dawes, C.J. 1998b. Biomass and photosynthetic responses to irradiance by a shallow and a deep water population of *Thalassia testudinum* on the west coast of Florida. *Bulletin of Marine Science* 62:89–96.
- Dawes, C. and J. Andorfer. 2002. Production of rhizome meristems by *Thalassia testudinum*. Pp. 185–197 in Greening, H.S. (ed.), *Proceedings, Seagrass Management: It's Not Just Nutrients!* Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Dawes, C.J., J. Andorfer, C. Rose, C. Uranowski, and N. Ehringer. 1997. Regrowth of the seagrass *Thalassia testudinum* into propeller scars. *Aquatic Botany* 59:139–155.
- Dawes, C.J., S.S. Bell, R.A. Davis, Jr., E.D. McCoy, H.R. Mushinsky, and J.L. Simon. 1995. Initial effects of Hurricane Andrew on the shoreline habitats of southwestern Florida. *Journal of Coastal Research* 21:103–110.
- Dawes, C.J., M. Chan, R. Chinn, E.W. Koch, A. Lazar, and D. Tomasko. 1987. Proximate composition, photosynthetic and respiratory responses of the seagrass *Halophila engelmannii* from Florida. *Aquatic Botany* 27:195–201.
- Dawes, C.J., M.O. Hall, and R.K. Reichert. 1985. Seasonal biomass and energy content in seagrass communities on the west coast of Florida. *Journal of Coastal Research* 1:255–262.
- Dawes, C.J. and J.M. Lawrence. 1979. Effects of blade removal on the proximate composition of the rhizome of the seagrass *Thalassia testudinum* Banks ex König. *Aquatic Botany* 7:255–266.
- Dawes, C.J. and J.M. Lawrence. 1980. Seasonal changes in the proximate constituents of the seagrasses *Thalassia testudinum*, *Halodule wrightii*, and *Syringodium filiforme*. Aquatic Botany 8:371–380.
- Dawes, C.J. and J.M. Lawrence. 1990. Seasonal changes in limestone and sand communities off the Florida west coast. *Pubblicazioni della Stazione zoologica di Napoli I: Marine Ecology* 11:97–104.

- Dawes, C.J., C.S. Lobban, and D.A. Tomasko. 1989. A comparison of the physiological ecology of the seagrasses *Halophila decipiens* Ostenfeld and *H. johnsonii* Eiseman from Florida. *Aquatic Botany* 33:149–154.
- Dawes, C.J. and D.A. Tomasko. 1988. Depth distribution of *Thalassia testudinum* in two meadows on the west coast of Florida USA: a difference in effect of light availability. *Pubblicazioni della Stazione zoologica di Napoli I: Marine Ecology* 9:123–130.
- Dawes, C.J. and J.F. van Breedveld. 1969. Benthic marine algae. *Memoirs of the Hourglass Cruises 1* (2). Florida Department of Natural Resources. St. Petersburg, Florida. 47 pp.
- Dawson, E.Y. 1966. *Marine Botany, An Introduction*. Holt, Rinehart, and Winston, Inc., New York, New York. 371 pp.
- den Hartog, C. 1970. *The Sea-Grasses of the World*. North Holland, Amsterdam, The Netherlands. 275 pp.
- Dennison, W.C., R.J. Orth, K.A. Moore, J.C. Stevenson, V. Carter, S. Kollar, P.W. Bergstrom, and R.A. Batiuk. 1993. Assessing water quality with submersed aquatic vegetation. *BioScience* 43:86–94.
- Dillon, K.S., J.P. Chanton, D.R. Corbett, and W.C. Burnett. 2002. Nitrogen isotopic compositions of seagrass and algae: implications for tracing nutrient sources in two Florida estuaries. Pp. 239–246 in Greening, H.S. (ed.), *Proceedings, Seagrass Management: It's Not Just Nutrients!* Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Dixon, L.K. 2000. Establishing light requirements for the seagrass *Thalassia testudinum*: an example from Tampa Bay, Florida. Pp. 9–31 *in* Bortone, S.A. (ed.), *Seagrasses: Monitoring, Ecology, Physiology, and Management.* CRC Press, Boca Raton, Florida. 318 pp.
- Dixon, L.K. 2002. Light requirements of Tampa Bay seagrasses: nutrient-related issues still pending. Pp. 21–28 in Greening, H.S. (ed.), Proceedings, Seagrass Management: It's Not Just Nutrients! Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Dixon, L.K. and G.J. Kirkpatrick. 1995. Causes of light attenuation with respect to seagrasses in upper and lower Charlotte Harbor. Final report to Southwest Florida Water Management District and Charlotte Harbor National Estuary Program. *Mote Marine Laboratory Technical Report 406*. Mote Marine Laboratory, Sarasota, Florida.
- Dixon, L.K. and G.J. Kirkpatrick. 1999. Causes of light attenuation with respect to seagrasses in upper

- and lower Charlotte Harbor. Final report to Southwest Florida Water Management District and Charlotte Harbor National Estuary Program. *Mote Marine Laboratory Technical Report 650*. Mote Marine Laboratory, Sarasota, Florida.
- Dixon, L.K. and J.R. Leverone. 1991. Evaluation of existing data on light requirements for the seagrasses *Thalassia testudinum* and *Halodule wrightii*. Final Report. *Mote Marine Laboratory Technical Report 291*. Mote Marine Laboratory, Sarasota, Florida. 15 pp.
- Dixon, L. K. and J.R. Leverone. 1995. Light requirements of *Thalassia testudinum* in Tampa Bay, Florida. Final Report. *Mote Marine Laboratory Technical Report 425*. Mote Marine Laboratory, Sarasota, Florida.
- Dixon, L. and J.R. Leverone. 1997. Annual light regime of light-limited *Thalassia testudinum* in Tampa Bay. Pp. 171–180 in Treat, S.F. (ed.), *Proceedings, Tampa Bay Area Scientific Information Symposium 3: Applying Our Knowledge*. Tampa Bay Regional Planning Council, Tampa, Florida. 396 pp.
- Doering, P.H. and R.H. Chamberlain. 2000. Experimental studies on salinity tolerance of turtle grass, *Thalassia testudinum*. Pp. 81–98 in Bortone, S.A. (ed.), *Seagrasses: Monitoring, Ecology, Physiology, and Management*. CRC Press, Boca Raton, Florida. 318 pp.
- Duarte, C.M. 1990. Seagrass nutrient content. *Marine Ecology Progress Series* 67:201–207.
- Duarte, C.M. 1991. Seagrass depth limits. *Aquatic Botany* 40:363–377.
- Duarte, C.M. 1995. Submerged aquatic vegetation in relation to different nutrient regimes. *Ophelia* 41:87–112.
- Duarte, C.M. 2002. The future of seagrass meadows. *Environmental Conservation* 29:192–206.
- Dunton, K.H. 1990. Production ecology of *Ruppia* maritima L. s.l. and *Halodule wrightii* Aschers. in two subtropical estuaries. *Journal of Experimental Marine Biology and Ecology* 143:147–164.
- Dunton, K.H. 1994. Seasonal growth and biomass of the subtropical seagrass *Halodule wrightii* in relation to continuous measurements of underwater irradiances. *Marine Biology* 120:479–489.
- Dunton, K.H. 1996. Photosynthetic production and biomass of the subtropical seagrass *Halodule wrightii* along an estuarine gradient. Estuaries 19:436–447.
- Durako, M.J. 1988. The seagrass bed. A community under assault. *Florida Naturalist* Fall 1998: 6–8.

- Durako, M.J. 1993. Photosynthetic utilization of CO_{2(aq)} and HCO₃- in *Thalassia testudinum* (Hydrocharitaceae). *Marine Biology* 115:373–380.
- Durako, M.J. 1994. Seagrass die-off in Florida Bay (USA): changes in shoot demographic characteristics and population dynamics in *Thalassia testudinum*. Marine Ecology Progress Series 110:59–66.
- Durako, M.J. and M.O. Hall. 1992. Effects of light on the stable carbon isotope composition of the seagrass *Thalassia testudinum*. *Marine Ecology Progress Series* 86:99–101.
- Durako, M.J., M.O. Hall, F. Sargent, and S. Peck. 1992. Propeller scars in seagrass beds: an assessment and experimental study of recolonization in Weedon Island State Preserve, Florida. Pp. 42–53 in Webb, F.J., Jr. (ed.), Proceedings of the Nineteenth Annual Conference on Wetlands Restoration and Creation. Hillsborough Community College, Tampa, Florida.
- Durako, M.J. and K.M. Kuss. 1994. Effects of Labyrinthula infection on the photosynthetic capacity of Thalassia testudinum. Bulletin of Marine Science 54:727–732.
- Durako, M.J. and M.D. Moffler. 1985. Observations on the reproductive ecology of *Thalassia testudinum* (Hydrocharitaceae). II. Leaf width as a secondary sex character. *Aquatic Botany* 21:265–275.
- Durako, M.J. and M.D. Moffler. 1987. Factors affecting the reproductive ecology of *Thalassia testudinum* (Hydrocharitaceae). *Aquatic Botany* 27:79–95.
- Durako, M.J. and W.M. Sackett. 1993. Effects of CO_{2(sq)} on the carbon isotopic composition of the seagrass *Thalassia testudinum* Banks ex König (Hydrocharitaceae). *Journal of Experimental Marine Biology and Ecology* 169:167–180.
- Eiseman, N.J. and C. McMillan. 1980. A new species of seagrass, *Halophila johnsonii*, from the Atlantic coast of Florida. *Aquatic Botany* 9:15–19.
- Eleuterius, L.N. 1987. Seagrass ecology along the coasts of Alabama, Louisiana, and Mississippi. Pp. 11–24 in Durako, M.J., R.C. Phillips, and R.R. Lewis III (eds.), Proceedings of the Symposium on Subtropical-Tropical Seagrasses of the Southeastern United States. Florida Marine Research Publications No. 42. Florida Marine Research Institute, St. Petersburg, Florida. 209 pp.
- Erskine, J.M. and M.S. Koch. 2000. Sulfide effects on *Thalassia testudinum* carbon balance and adenylate energy charge. *Aquatic Botany* 67:275–285.
- Estevez, E.D. 2000. Matching salinity metrics to estu-

- arine seagrasses for freshwater inflow management. Pp. 295–307 in Bortone, S.A. (ed.), Seagrasses: Monitoring, Ecology, Physiology, and Management. CRC Press, Boca Raton, Florida. 318 pp.
- Ferguson, R.L., B.T. Pawlak, and L.L. Wood. 1993. Flowering of the seagrass *Halodule wrightii* in North Carolina, USA. *Aquatic Botany* 46: 91–98.
- Fishman, J.R. and R.J. Orth. 1996. Effects of predation on Zostera marina L. seed abundance. Journal of Experimental Marine Biology and Ecology 198:11–26.
- Fitzpatrick, J. and H. Kirkman. 1995. Effects of prolonged shading stress on growth and survival of seagrass *Posidonia australis* in Jervis Bay, New South Wales, Australia. *Marine Ecology Progress Series* 127:279–289.
- Fletcher, S.W. and W.W. Fletcher. 1995. Factors affecting changes in seagrass distribution and diversity patterns in the Indian River Lagoon complex between 1940 and 1992. *Bulletin of Marine Science* 57:49–58.
- Florida Institute of Oceanography. 1994. The West Florida Shelf: past, present and future. Final Workshop Report. Florida Institute of Oceanography, St. Petersburg, Florida.
- Folit, R. and J. Morris. 1992. Beds, boats, and buoys: a study in protecting seagrass beds from motorboat propeller damage: Early Action Demonstration Project, Sarasota Bay National Estuary Program, Sarasota, Florida.
- Fonseca, M.S. 1989. Sediment stabilization by *Halophila decipiens* in comparison to other seagrasses. *Estuarine*, *Coastal and Shelf Science* 29:501–507.
- Fonseca, M.S. and S.S. Bell. 1998. Influence of physical setting on seagrass landscapes near Beaufort, North Carolina, USA. *Marine Ecology Progress Series* 171:109–121.
- Fonseca, M.S., S.S. Bell, W.J. Kenworthy, P. Whitfield, M. Finkbeiner, and M.O. Hall. 2001. An examination of scale dependence in the assessment of temporally and spatially dynamic deepwater seagrass (*Halophila decipiens*) beds of the West Florida Shelf. *Abstract* in Estuarine Research Federation Meetings, November 2001. St. Petersburg, Florida.
- Fonseca, M.S., W.J. Kenworthy, and F.X. Courtney. 1996a. Development of planted seagrass beds in Tampa Bay, Florida, USA. I. Plant components. *Marine Ecology Progress Series* 132:127–139.
- Fonseca, M.S., W.J. Kenworthy, and G.W. Thayer.

- 1998. Guidelines for the conservation and restoration of seagrasses in the United States and adjacent waters. National Oceanic and Atmospheric Administration Coastal Ocean Program Decision Analysis Series, No. 12. National Oceanic and Atmospheric Administration Coastal Ocean Office, Silver Spring, Maryland. 222 pp.
- Fonseca, M.S., D.L. Meyer, and M.O. Hall. 1996b. Development of planted seagrass beds in Tampa Bay, Florida, USA. II. Faunal components. *Marine Ecology Progress Series* 132:141–156.
- Fonseca, M.S., B.D. Robbins, P.E. Whitfield, L. Woods, and P. Clinton. 2002. Evaluating the effect of offshore sandbars on seagrass recovery and restoration in Tampa Bay through ecological forecasting and hind-casting of exposure to waves. *Tampa Bay Estuary Program Technical Publication #07-02*. Tampa Bay Estuary Program, St. Petersburg, Florida. 48 pp.
- Fourqurean, J.W., G.V.N. Powell, W.J. Kenworthy, and J.C. Zieman. 1995. The effects of long-term manipulation of nutrient supply on competition between the seagrasses *Thalassia testudinum* and *Halodule wrightii* in Florida Bay. *Oikos* 72:349–358.
- Fourqurean, J.W. and M.B. Robblee. 1999. Florida Bay: a brief history of recent ecological changes. *Estuaries* 22:345–357.
- Fourqurean, J.W. and J.C. Zieman. 1991. Photosynthesis, respiration and whole plant carbon budget of the seagrass *Thalassia testudinum*. *Marine Ecology Progress Series* 69:161–170.
- Fourqurean, J.W., J.C. Zieman, and G.V.N. Powell. 1992. Relationships between porewater nutrients and seagrasses in a subtropical carbonate environment. *Marine Biology* 114:57–65.
- Frankovich, T.A. and J.W. Fourqurean. 1997. Seagrass epiphyte loads along a nutrient availability gradient, Florida Bay, USA. *Marine Ecology Progress Series* 159:37–50.
- Fry, B. 1981. Tracing shrimp migrations and diets using natural variations in stable isotopes. Ph.D. Dissertation. University of Texas, Austin, Texas. 225 pp.
- Fry, B. 1984. ¹³C/¹²C ratios and the trophic importance of algae in Florida *Syringodium filiforme* seagrass meadows. *Marine Biology* 79:11–19.
- Fry, B., S.A. Macko, and J.C. Zieman. 1987. Review of stable isotopic investigations of food webs in seagrass meadows. Pp. 189–209 in Durako, M.J., R.C. Phillips, and R.R. Lewis III. (eds.), Proceedings of the Symposium on Subtropical-Tropical Seagrasses of

- the Southeastern United States. Florida Marine Research Publications No. 42. Florida Marine Research Institute, St. Petersburg, Florida. 209 pp.
- Fry, B. and P.L. Parker. 1979. Animal diet in Texas seagrass meadows: δ¹³C evidence for the importance of benthic plants. *Estuarine and Coastal Marine Science* 8:499–509.
- Fry, B. and R.W. Virnstein. 1988. Leaf production and export of the seagrass *Syringodium filiforme* Kütz. in Indian River Lagoon, Florida. *Aquatic Botany* 30:261–266.
- Gacia, E. 1999. Leaf dynamics and shoot production of the seagrass *Thalassia testudinum* in the Indian River Lagoon (Florida). *Botanica Marina* 42:97–102.
- Gacia, E., M.M. Littler, and D.S. Littler. 1999. An experimental test of the capacity of food web interactions (fish-epiphytes-seagrasses) to offset the negative consequences of eutrophication on seagrass communities. *Estuarine, Coastal and Shelf Science* 48:757–766.
- Gallegos, C.L. and W.J. Kenworthy. 1996. Seagrass depth limits in the Indian River Lagoon (Florida, U.S.A.): application of an optical water quality model. Estuarine, Coastal and Shelf Science 42:267–288.
- Gessner, F. 1970. Temperature: plants. Pp. 363–406 in Kinne, O. (ed.), Marine Ecology: A Comprehensive, Integrated Treatise on Life in Oceans and Coastal Waters. Volume 1. Environmental Factors. Part 1. Wiley-Interscience, London, England. 681 pp.
- Gilmore, R.G. 1987. Subtropical-tropical seagrass communities of the southeastern United States: fishes and fish communities. Pp. 117–137 in Durako, M.J., R.C. Phillips, and R.R. Lewis III (eds.), Proceedings of the Symposium on Subtropical-Tropical Seagrasses of the Southeastern United States. Florida Marine Research Publications No. 42. Florida Marine Research Institute, St. Petersburg, Florida. 209 pp.
- Godcharles, M.F. 1971. A study of the effects of a commercial hydraulic clam dredge on benthic communities in estuarine areas. *Florida Marine Research Laboratory Technical Series 64*. Florida Department of Natural Resources, St. Petersburg, Florida. 51 pp.
- Grabe, S. 1999. Sediment quality. Pp.1–6 in Pribble, J.R., A.J. Janicki, and H. Greening (eds.), Baywide environmental monitoring report, 1993–1998, Tampa Bay, Florida. Tampa Bay Estuary Program Technical Publication # 07-99. Tampa Bay Estuary Program, St. Petersburg, Florida.

- Greening, H. S. (ed.). 2002a. *Proceedings, Seagrass Management: It's Not Just Nutrients!* Proceedings of a symposium, August 22–24, 2000, St. Petersburg, Florida. Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Greening, H. S. 2002b. Implementing the Tampa Bay seagrass restoration management strategy. Pp. 29–37 in Greening, H.S. (ed.), Proceedings, Seagrass Management: It's Not Just Nutrients! Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Greening, H.S. and R.J. Livingston. 1982. Diel variation in the structure of seagrass-associated epibenthic macroinvertebrate communities. *Marine Ecology Progress Series* 7:147–156.
- Greenway, M. 1995. Trophic relationships of macrofauna within a Jamaican seagrass meadow and the role of the echinoid *Lytechinus variegatus* (Lamarck). *Bulletin of Marine Science* 56:719–736.
- Gulf of Mexico Program (GMP). 2003. Gulf of Mexico Program objectives, sub-objectives, and annual performance goals for fiscal year 2003. Stennis Space Center, Mississippi.
- Haddad, K.D. 1989. Habitat trends and fisheries in Tampa and Sarasota bays. In Estevez, E.D. (ed.), Tampa and Sarasota Bays: Issues, Resources, and Status and Management. National Oceanic and Atmospheric Administration Estuary-of-the-Month Seminar Series No. 11. National Oceanic and Atmospheric Administration, Washington, D.C.
- Hall, M.O. 1988. Dynamics and interactions of epiphytic macroalgae and meiofauna on the seagrass *Thalassia testudinum*. Ph.D. Dissertation, University of South Florida, Tampa, Florida.
- Hall, M.O. and S.S. Bell. 1988. Response of small motile epifauna to complexity of epiphytic algae on seagrass blades. *Journal of Marine Research* 46:613–630.
- Hall, M.O. and S.S. Bell. 1993. Meiofauna on the seagrass *Thalassia testudinum*: population characteristics of harpacticoid copepods and associations with algal epiphytes. *Marine Biology* 116:137–146.
- Hall, M.O., M.J. Durako, J.W. Fourqurean, and J.C. Zieman. 1999. Decadal-scale changes in seagrass distribution and abundance in Florida Bay. *Estuaries* 22:445–459.
- Hamm, D. and H. Humm. 1976. Benthic algae of the Anclote Estuary. II. Bottom-dwelling species. *Florida Scientist* 39:209–229.
- Harper, J.L. 1990. *Population Biology of Plants*, 8th Ed. Academic Press, London, England. 892 pp.

- Harris, F.L. 1997. The effects of epiphytic algae on blade growth of *Thalassia testudinum* Banks ex König in Sarasota Bay, Florida. M.S. Thesis. University of South Florida, Tampa, Florida.
- Harris, B.A., K.D. Haddad, K.A. Steidinger, and J.A. Huff. 1983. Assessment of fisheries habitat: Charlotte Harbor and Lake Worth, Florida. Final Report. Florida Department of Natural Resources. Available from the Florida Fish and Wildlife Conservation Commission, Florida Marine Research Institute, St. Petersburg, Florida. 227 pp.
- Hauxwell, J., J. Cebrián, J.A. Herrera-Silveira, J. Ramírez-Ramírez, A. Zaldivar-Jimenez, N. Gomez, and N. Aranda-Cirerol. 2001. Measuring production of *Halodule wrightii*: additional evidence suggests clipping underestimates growth rate. *Aquatic Botany* 69:41–54.
- Heck, K.L., Jr., G. Hays, and R.J. Orth. 2003. Critical evaluation of the nursery role hypothesis for seagrass meadows. *Marine Ecology Progress Series* 253:123–136.
- Heck, K.L., Jr. and E.D. McCoy. 1979. Biogeography of seagrasses: evidence from associated organisms. Proceedings of the First International Symposium on Marine Biogeography and Evolution in the Southern Hemisphere. New Zealand Department of Science and Industrial Research Information Series 137:109–127.
- Heck, K.L., Jr., and J.F. Valentine. 1995. Sea urchin herbivory: evidence for long-lasting effects in subtropical seagrass meadows. *Journal of Experimental Marine Biology and Ecology* 189:205–217.
- Hemminga, M.A. and C.M. Duarte. 2000. *Seagrass Ecology*. Cambridge University Press, Cambridge, Massachusetts. 298 pp.
- Hemminga, M.A. and M.A. Mateo. 1996. Stable carbon isotopes in seagrasses: variability in ratios and use in ecological studies. *Marine Ecology Progress Series* 140:285–298.
- Hillman, K., D.I. Walker, A.W.D. Larkum, and A.J. McComb. 1989. Productivity and nutrient limitation. Pp. 635–685 in Larkum, A.W.D., A.J. McComb, and S.A. Shepherd (eds.), Biology of Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region. Elsevier, Amsterdam, The Netherlands. 841 pp.
- Holmquist, J.G. 1994. Benthic macroalgae as a dispersal mechanism for fauna: influence of a marine tumbleweed. *Journal of Experimental Marine Biology and Ecology* 180:235–251.

- Holmquist, J.G., G.V.N. Powell, and S.M. Sogard. 1989. Decapod and stomatopod assemblages on a system of seagrass-covered mud banks in Florida Bay. *Marine Biology* 100:473–483.
- Howard, R.K. and F.T. Short. 1986. Seagrass growth and survivorship under the influence of epiphyte grazers. *Aquatic Botany* 24:287–302.
- Iverson, R.L. and H.F. Bittaker. 1986. Seagrass distribution and abundance in eastern Gulf of Mexico coastal waters. *Estuarine, Coastal and Shelf Science* 22:577–602.
- Jaap, W.C. and P. Hallock. 1990. Coral reefs. Pp. 574–616 in Myers, M.L. and J.J. Ewel (eds.), *Ecosystems of Florida*. University of Central Florida Press, Orlando, Florida. 765 pp.
- Jagels, R. and A. Barnabas. 1989. Variation in leaf ultrastructure of *Ruppia maritima* L. along a salinity gradient. *Aquatic Botany* 33:207–221.
- Janicki, A.J., D.L. Wade, and D.E. Robison. 1994. Habitat protection and restoration targets for Tampa Bay. Tampa Bay National Estuary Program Technical Publication #07-93. Tampa Bay National Estuary Program. St. Petersburg, Florida.
- Janicki, A., D.L. Wade, and J.R. Pribble. 1999. Development of a process to track the status of chlorophyll and light attenuation to support seagrass restoration goals in Tampa Bay. Janicki Environmental, Inc., St. Petersburg, Florida.
- Jensen, M.L. and N. Nakai. 1961. Sources and isotopic composition of atmospheric sulfur. *Science* 134: 2102–2104.
- Jensen, S. and S. Bell. 2001. Seagrass growth and patch dynamics: cross-scale morphological plasticity. *Plant Ecology* 155:201–217.
- Johansson, J.O.R. 1991. Long-term trends of nitrogen loading, water quality and biological indicators in Hillsborough Bay, Florida. Pp.157–176 in Treat, S.F. and P.A. Clark (eds.), Proceedings, Tampa Bay Area Scientific Information Symposium 2: The Watershed. TEXT, Tampa, Florida. 528 pp.
- Johansson, J.O.R. 2002a. Historical overview of Tampa Bay water quality and seagrass: issues and trends. Pp.1–10 in Greening, H.S. (ed.), *Proceedings, Sea*grass Management: It's Not Just Nutrients! Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Johansson, J.O.R. 2002b. Water depth (MTL) at the deep edge of seagrass meadows in Tampa Bay measured by GPS carrier-phase processing: evaluation of the technique. Pp.151–168 in Greening, H.S. (ed.), Proceedings, Seagrass Management: It's Not

Just Nutrients! Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.

Johansson, J.O.R. and H.S. Greening. 2000. Seagrass restoration in Tampa Bay: a resource-based approach to estuarine management. Pp. 279–293 in Bortone, S.A. (ed.), Seagrasses: Monitoring, Ecology, Physiology, and Management. CRC Press, Boca Raton, Florida. 318 pp.

Johansson, J.O.R. and T. Ries. 1997. Seagrass in Tampa Bay: historic trends and future expectations. Pp. 139–150 in Treat, S.F. (ed.), Proceedings, Tampa Bay Area Scientific Information Symposium 3: Applying Our Knowledge. Tampa Bay Estuary Program, St. Petersburg, Florida. 396 pp.

Jones, J.A. 1968. Primary productivity by the tropical marine turtle grass *Thalassia testudinum* König, and its epiphytes. Ph.D. Dissertation, University of Miami, Miami, Florida. 196 pp.

Josselyn, M., M. Fonseca, T. Niesen, and R. Larson. 1986. Biomass, production and decomposition of a deep water seagrass, *Halophila decipiens* Ostenf. *Aquatic Botany* 25:47–61.

Kaldy, J.E., N. Fowler, and K.H. Dunton. 1999. Critical assessment of *Thalassia testudinum* (turtle grass) aging techniques: implications for demographic inferences. *Marine Ecology Progress Series* 181:279–288.

Katz, B.G., R.S. DeHan, J.J. Hirten, and J.S. Catches. 1997. Interactions between ground water and surface water in the Suwannee River basin, Florida. *Journal of the American Water Resources Association* 33:1237–1254.

Kelly, P.O. 1995. Long-term trends of macroalgae in Hillsborough Bay. *Florida Scientist* 58:179–191.

Kenworthy, W.J. 1993. Defining the ecological light compensation point of seagrasses in the Indian River Lagoon. Pp. 195–210 in Morris, L.J. and D.A. Tomasko (eds.), Proceedings and Conclusions of Workshops on Submerged Aquatic Vegetation Initiative and Photosynthetically Active Radiation. St. John's River Water Management District, Palatka, Florida.

Kenworthy, W.J., C.A. Currin, M.S. Fonseca, and G. Smith. 1989. Production, decomposition, and heterotrophic utilization of the seagrass *Halophila decipiens* in a submarine canyon. *Marine Ecology Progress Series* 51:277–290.

Kenworthy, W.J., C. Currin, G. Smith, and G. Thayer. 1987. The abundance, biomass and acetylene reduction activity of bacteria associated with decomposing rhizomes of two seagrasses, Zostera marina and Thalassia testudinum, Aquatic Botany 27:97–119.

Kenworthy, W.J. and M.S. Fonseca. 1996. Light requirements of seagrasses *Halodule wrightii* and *Syringodium filiforme* derived from the relationship between diffuse light attenuation and maximum depth distribution. *Estuaries* 19:740–750.

Kenworthy, W.J., M.S. Fonseca, and G.W. Thayer. 1988a. A comparison of wind-wave and boat wake energy in Hobe Sound: implications for seagrass growth. Annual Report to the U.S. Fish and Wildlife Service Sirenia Project, Gainesville, Florida. 21 pp.

Kenworthy, W.J. and E.E. Haunert (eds.). 1991. The Light Requirements of Seagrasses: Proceedings of a Workshop to Examine the Capability of Water Quality Criteria, Standards and Monitoring Programs to Protect Seagrasses. National Oceanic and Atmospheric Administration Technical Memorandum NMFS-SEFC-287. National Oceanographic and Atmospheric Association, Washington, D.C. 187 pp.

Kenworthy, W.J., G.W. Thayer, and M.S. Fonseca. 1988b. The utilization of seagrass meadows by fishery organisms. Pp. 548–560 in Hook, D.D., W.H. McKee, Jr., H.K. Smith, J. Gregory, V.G. Burrell, Jr., M.R. Devoe, R.E. Sojka, S. Gilbert, R. Banks, L.H. Stolzy, C. Brooks, T.D. Matthews, and T.H. Shear (eds.), The Ecology and Management of Wetlands. Volume 1. Ecology of Wetlands. Timber Press, Portland, Oregon. 592 pp.

Kharlamenko, V.I., S.I. Kiyashko, A.B. Imbs, and D.I. Vyshkvartzev. 2001. Identification of food sources of invertebrates from the seagrass *Zostera marina* community using carbon and sulfur stable isotope ratio and fatty acid analyses. *Marine Ecology Progress Series* 220:103–117.

Kikuchi, T. 1980. Faunal relationships in temperate seagrass beds. Pp. 153–172 in Phillips, R.C. and C.P. McRoy (eds.), Handbook of Seagrass Biology—An Ecosystem Perspective. Garland STPM Press, New York, New York. 353 pp.

Kikuchi, T. and J.P. Peres. 1977. Consumer ecology of seagrass beds. Chapter 5 in McRoy, C.P. and C. Helfferich (eds.), Seagrass Ecosystems: A Scientific Perspective. Marcel Dekker, New York, New York. 314 pp.

Kirsten, J.H., C.J. Dawes, and B.J. Cochrane. 1998. Randomly amplified polymorphism detection (RAPD) reveals high genetic diversity in *Thalassia testudinum* Banks ex König (Turtlegrass). *Aquatic Botany* 61:269–287.

Klumpp, D.W., R.K. Howard, and D.A. Pollard. 1989.

- Trophodynamics and nutritional ecology of seagrass communities. Pp. 394–457 in Larkum, A.W.D., A.J. McComb, and S.A. Shepherd (eds.), Biology of Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region. Elsevier, Amsterdam, The Netherlands. 841 pp.
- Knowles, L.L. and S.S. Bell. 1998. The influence of habitat structure in faunal-habitat associations in a Tampa Bay seagrass system, Florida. *Bulletin of Marine Science* 62:781–794.
- Koch, E.W. 1999a. Preliminary evidence on the interdependent effect of currents and porewater geochemistry on *Thalassia testudinum* Banks ex König seedlings. *Aquatic Botany* 63:95–102.
- Koch, E.W. 1999b. Sediment resuspension in a shallow *Thalassia testudinum* Banks ex König bed. *Aquatic Botany* 65:269–280.
- Koch, E.W. 2001. Beyond light: physical, geological, and geochemical parameters as possible submersed aquatic vegetation habitat requirements. *Estuaries* 24:1–17.
- Koch, E.W. and S. Beer. 1996. Tides, light and the distribution of *Zostera marina* in Long Island Sound, USA. *Aquatic Botany* 53:97–107.
- Koch, E.W. and C. J. Dawes. 1991a. Influence of salinity and temperature on the germination of *Ruppia maritima* L. from the north Atlantic and Gulf of Mexico. *Aquatic Botany* 40:387–391.
- Koch, E.W. and C.J. Dawes. 1991b. Ecotypic differentiation in populations of *Ruppia maritima* L. germinated from seeds and cultured under algaefree laboratory conditions. *Journal of Experimental Marine Biology and Ecology* 152:145–159.
- Koch, E.W. and G. Gust. 1999. Water flow in tide- and wave-dominated beds of the seagrass *Thalassia testudinum*. *Marine Ecology Progress Series* 184:63–72.
- Koch, M.S. and J.M. Erskine. 2001. Sulfide as a phytotoxin to the tropical seagrass *Thalassia testudinum*: interactions with light, salinity, and temperature. *Journal of Experimental Marine Biology and Ecology* 266:81–95.
- Kraemer, G.P. and M.D. Hanisak. 2000. Physiological and growth responses of *Thalassia testudinum* to environmentally relevant periods of low irradiance. *Aquatic Botany* 67:287–300.
- Kruzcynski, A. 1994. Anatomical and cytological adaptations to salinity in the leaves of *Ruppia maritima*. M.S. Thesis. University of South Florida, Tampa, Florida.
- Kruer, C.R. 1994. Mapping assessment of vessel dam-

- age to shallow seagrasses in the Florida Keys. Final Report for contract 47-10-123-L3. Florida Department of Natural Resources and Florida Institute of Oceanography, St. Petersburg, Florida.
- Kuo, J. and C. den Hartog. 1989. Evolution and biogeography of seagrasses. Pp.112–156 in Larkum, A.W.D., A.J. McComb, and S.A. Shepherd (eds.), Biology of Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region. Elsevier, Amsterdam, The Netherlands. 841 pp.
- Kuo, J. and C. den Hartog. 2001. Seagrass taxonomy and identification key. Pp. 31–58 *in* Short, F.T. and R.G. Coles (eds.), *Global Seagrass Research Methods*. Elsevier Science B.V., Amsterdam, The Netherlands. 473 pp.
- Kuo, J. and A.J. McComb. 1989. Seagrass taxonomy, structure and development. Pp. 6–73 in Larkum, A.W.D., A.J. McComb, and S.A. Shepherd (eds.), Biology of Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region. Elsevier, Amsterdam, The Netherlands. 841 pp.
- Kurz, R.C., D.A. Tomasko, D. Burdick, T. Ries, K. Patterson, and R. Finck. 1999. Summary of recent trends in seagrass distributions in southwest Florida coastal waters. Surface Water Improvement and Management (SWIM) Program Technical Report. Southwest Florida Water Management District, Brooksville, Florida.
- Kurz, R.C., D.A. Tomasko, D. Burdick, T.F. Ries, S.K. Patterson, and R. Finck. 2000. Recent trends in seagrass distributions in southwest Florida coastal waters. Pp.157–166 in Bortone, S.A. (ed.), Seagrasses: Monitoring, Ecology, Physiology, and Management. CRC Press, Boca Raton, Florida. 318 pp.
- Lapointe, B.E. and P.A. Clark. 1992. Nutrient inputs from the watershed and coastal eutrophication in the Florida Keys. *Estuaries* 15:465–476.
- Lapointe, B.E., D.A. Tomasko, and W.R. Matzie. 1994. Eutrophication and trophic state classification of seagrass communities in the Florida Keys. *Bulletin of Marine Science* 54:696–717.
- Larkum, A.W.D., G. Roberts, J. Kuo, and S. Strother. 1989. Gaseous movement in seagrasses. Pp. 686–722 in Larkum, A.W.D., A.J. McComb, and S.A. Shepherd (eds.), Biology of Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region. Elsevier, Amsterdam, The Netherlands. 841 pp.
- Larkum, A.W.D. and W.F. Wood. 1993. The effect of UV-B radiation on photosynthesis and respiration

- of phytoplankton, benthic macroalgae and seagrasses. *Photosynthesis Research* 36:17–23.
- Lazar, A.C. and C.J. Dawes. 1991. A seasonal study of the seagrass *Ruppia maritima* L. in Tampa Bay, Florida. Organic constituents and tolerances to salinity and temperature. *Botanica Marina* 34:265–269.
- Lee, K.-S. and K.H. Dunton. 1997. Effects of *in situ* light reduction on the maintenance, growth and partitioning of the carbon resources in *Thalassia testudinum* Banks ex König. *Journal of Experimental Marine Biology and Ecology* 210:53–73.
- Lee, K.-S. and K.H. Dunton. 2000a. Diurnal changes in pore water sulfide concentrations in the seagrass *Thalassia testudinum* beds: the effects of seagrasses on sulfide dynamics. *Journal of Experimental Marine Biology and Ecology* 255:201–214.
- Lee, K.-S. and K.H. Dunton. 2000b. Effects of nitrogen enrichment on biomass allocation, growth, and leaf morphology of the seagrass *Thalassia testudinum*. *Marine Ecology Progress Series* 196:39–48.
- Lefebvre, L.W., T.J. O'Shea, G.B. Rathbun, and R.C. Best. 1989. Distribution, status, and biogeography of the West Indian Manatee. Pp. 567–610 in Woods, C.A. (ed.), Biogeography of the West Indies: Past, Present, and Future. Sandhill Crane Press, Gainesville, Florida. 878 pp.
- Lewis, F.G., III. 1984. Distribution of macrobenthic crustaceans associated with *Thalassia*, *Halodule*, and bare sand substrata. *Marine Ecology Progress Series* 19:101–113.
- Lewis, F.G., III. 1987. Crustacean epifauna of seagrass and macroalgae in Apalachee Bay, Florida, USA. *Marine Biology* 94:219–229.
- Lewis, F.G., III and A.W. Stoner. 1981. An examination of methods for sampling macrobenthos in seagrass meadows. *Bulletin of Marine Science* 31:116–124.
- Lewis, F.G., III and A.W. Stoner. 1983. Distribution of macrofauna within seagrass beds: an explanation for patterns of abundance. *Bulletin of Marine Science* 33: 296–304.
- Lewis, R.R., III. 2002. The potential importance of the longshore bar system to the persistence and restoration of Tampa Bay seagrass meadows. Pp.177–183 in Greening, H.S. (ed.), *Proceedings, Seagrass Management: It's Not Just Nutrients!* Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Lewis, R.R., III, P. Clark, W.K. Fehring, H.S. Greening, R. Johansson, and R.T. Paul. 1998. The reha-

- bilitation of the Tampa Bay estuary, Florida, USA: an example of successful integrated coastal management. *Marine Pollution Bulletin* 37:468–473.
- Lewis, R.R., III., M.J. Durako, M.D. Moffler, and R.C. Phillips. 1985a. Seagrass meadows of Tampa Bay: a review. Pp. 210–246 in Treat, S.A.F., J.L. Simon, R.R. Lewis III, and R.L Whitman, Jr. (eds.), Proceedings, Tampa Bay Area Scientific Information Symposium. Bellweather Press, Tampa, Florida. 663 pp.
- Lewis, R.R., III, R.G. Gilmore, Jr., D.W. Crewz, and W.E. Odum. 1985b. Mangrove habitat and fishery resources of Florida. Pp. 281–336 in Seaman, W.E., Jr. (ed.), Florida Aquatic Habitat and Fishery Resources. Florida Chapter of American Fisheries Society, Kissimmee, Florida. 543 pp.
- Lewis, R.R., III, K.D. Haddad, and J.O.R. Johansson. 1991. Recent aerial expansion of seagrass meadows in Tampa Bay, Florida: real bay improvement or drought-induced? Pp. 189–192 in Treat, S.F. and P.A. Clark (eds.), Proceedings: Tampa Bay Area Scientific Information Symposium 2: The Watershed. TEXT, Tampa, Florida. 528 pp.
- Lewis, R.R., III, A.B. Hodgson, M. Tooze, and C.D. Kruer. 2002. The distribution of seagrass and benthic habitats westward of the patch reef system boundary in Biscayne National Park, Florida, USA. Pp. 125–132 in Greening, H.S. (ed.), *Proceedings, Seagrass Management: It's Not Just Nutrients!* Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Lewis, R.R., III and R.C. Phillips. 1980. Occurrence of seeds and seedlings of *Thalassia testudinum* Banks ex König in the Florida Keys (U.S.A.). *Aquatic Botany* 9:377–380.
- Lin, G., T. Banks, and L.O. Sternberg. 1991. Variation in δ^{13} C values for the seagrass *Thalassia testudinum* and its relations to mangrove carbon. *Aquatic Botany* 40:333–341.
- Littler, D.S. and M.M. Littler. 2000. Caribbean Reef Plants. Offshore Graphics Inc., P.O. Box 6139, Washington, D.C., 20044.
- Livingston, R.J. 1987. Historic trends of human impacts on seagrass meadows in Florida. Pp.139–151 in Durako, M.J., R.C. Phillips, and R.R. Lewis III. (eds.), Proceedings of the Symposium on Subtropical-Tropical Seagrasses of the Southeastern United States. Florida Marine Research Publications No. 42. Florida Marine Research Institute, St. Petersburg, Florida. 209 pp.
- Livingston, R.J. 1990. Inshore marine habitats. Pp. 549–573 *in* Myers, R.L. and J.J. Ewel (eds.), *Ecosys*-

- tems of Florida. University of Central Florida Press, Orlando, Florida. 765 pp.
- Livingston, R.J. 1993. River/Gulf study. Chapter III. Light transmission characteristics and submerged aquatic vegetation. Report prepared by Environmental Planning and Analysis for the Buckeye Kraft L.P., Perry, Florida.
- Livingston, R.J., S.E. McGlynn, and X. Niu. 1998. Factors controlling seagrass growth in a Gulf coastal system: water and sediment quality and light. *Aquatic Botany* 60:135–159.
- Lobban, C.S. and P.J. Harrison. 1994. Seaweed Ecology and Physiology. Cambridge University Press, Cambridge, Massachusetts. 366 pp.
- Loflin, R.K. 1995. The effects of docks on seagrass beds in the Charlotte Harbor estuary. *Florida Scientist* 58:198–205.
- Lores, E.M., E. Pasko, J.M. Patrick, R.L. Quarles, J. Campbell, and J. Macauley. 2000. Mapping and monitoring of submerged aquatic vegetation in Escambia-Pensacola Bay system, Florida. *Gulf of Mexico Science* 2000:1–14.
- Lumbert, S.H., C. den Hartog, R.C. Phillips, and F.S. Olsen. 1984. The occurrence of fossil seagrasses in the Avon Park Formation (late Middle Eocene), Levy County, Florida (U.S.A.). Aquatic Botany 20:121–129.
- Macauley, J.M., J.R. Clark, and W.A. Price. 1988. Seasonal changes in the standing crop and chlorophyll content of *Thalassia testudinum* Banks ex König and its epiphytes in the northern Gulf of Mexico. *Aquatic Botany* 31:277–287.
- Maciá, S. 1999. The effects of sea urchin grazing and drift algal blooms on a subtropical seagrass bed community. *Journal of Experimental Marine Biology and Ecology* 246:53–47.
- Maciá, S. and D. Lirman. 1999. Destruction of Florida Bay seagrass by a grazing front of sea urchins. *Bulletin of Marine Science* 65:593–601.
- Macko, S.A. 1981. Stable nitrogen isotope ratios as tracers of organic geochemical processes. Ph.D. Dissertation, University of Texas, Austin, Texas. 194 pp.
- Madley, K.A., J.M. Lewis, J.D. Krolick, and E.C. Santi. 2003. Florida Marine Research Institute Website Seagrass Pages. http://www.floridamarine.org/features/category sub.asp?id=1306.
- Mahadevan, S. and G.W. Patton. 1979. A study of sieve (screen-mesh opening) size effects on benthic fauna collected from Anclote Anchorage. *Environmental*

- Protection Agency Technical Report #68-01-5016. Washington, DC.
- Marbá, N. and C.M. Duarte. 1998. Rhizome elongation and seagrass clonal growth. *Marine Ecology Progress Series* 174:269–280.
- Marbá, N., M.E. Gallegos, M. Merino, and C.M. Duarte. 1994. Vertical growth of *Thalassia testudinum:* seasonal and interannual variability. *Aguatic Botany* 47:1–11.
- Marx, J.M. and W.F. Herrnkind. 1985. Macroalgae (Rhodophyta: *Laurencia* spp.) as habitat for young juvenile spiny lobsters, *Panulirus argus. Bulletin of Marine Science* 36:423–431.
- Matthews, T.R., A.C. Lazar, and J.H. Hunt. 1991. Aerial observations of boating impacts to shallow water seagrass beds in the Florida Keys. Report to the Florida Department of Natural Resources, Office of Marine Programs and Planning, Tallahassee, Florida. 11 pp.
- Mattson, R.A. 1995. Submerged macrophyte communities. Chapter 6 in A Data Collection Program for Selected Coastal Estuaries in Hernando, Citrus and Levy Counties, Florida. Southwest Florida Water Management District, Brooksville, Florida.
- Mattson, R.A. 2000. Seagrass ecosystem characteristics and research and management needs in the Florida Big Bend. Pp. 259–277 in Bortone, S.A. (ed.), Seagrasses: Monitoring, Ecology, Physiology, and Management. CRC Press, Boca Raton, Florida. 318 pp.
- McMichael, R.H., Jr. and K.M. Peters. 1989. Early life history of spotted sea trout, *Cynoscion nebulosus* (Pisces: Sciaenidae), in Tampa Bay. *Estuaries* 12:98–110.
- McMillan, C. 1978. Morphogeographic variation under controlled conditions in five seagrasses, *Thalassia testudinum*, *Halodule wrightii*, *Syringodium filiforme*, *Halophila engelmannii*, and *Zostera marina*. Aquatic Botany 4:169–189.
- McMillan, C. 1979. Differentiation in response to chilling temperatures among populations of three marine spermatophytes, *Thalassia testudinum*, *Syringodium filiforme*, and *Halodule wrightii*. *American Journal of Botany* 66:810–819.
- McMillan, C. 1980. Isozymes of tropical seagrasses from the Indo-Pacific and the Gulf of Mexico-Caribbean. *Aquatic Botany* 8:163–172.
- McMillan, C. 1982. Reproductive physiology of tropical seagrasses. *Aquatic Botany* 14:245–258.
- McMillan, C. 1984. The distribution of tropical sea-

- grasses with relation to their tolerance of high temperatures. *Aquatic Botany* 19:369–379.
- McNulty, J.K., W.N. Lindall, and J.E. Sykes. 1972. Cooperative Gulf of Mexico estuarine inventory and study, Florida. Phase I: Area description. National Oceanic and Atmospheric Administration Technical Report, *National Marine Fisheries Service Circular No. 368*. National Oceanic and Atmospheric Administration, Washington, D.C.
- McPherson, B. F. and R.L. Miller. 1994. Causes of light attenuation in Tampa Bay and Charlotte Harbor, southwestern Florida. *Water Resources Bulletin* 30:43–53.
- McRoy, C.P. and C. Helfferich. 1980. Applied aspects of seagrasses. Pp. 297–342 in Phillips, R.C. and C.P. McRoy (eds.), *Handbook of Seagrass Biology—An Ecosystem Perspective*. Garland STPM Press, New York. 353 pp.
- McRoy, C.P. and C. McMillan 1977. Production ecology and physiology of seagrasses. Pp. 53–81 in McRoy, C.P and C. Helfferich (eds.), Seagrass Ecosystems: A Scientific Perspective. Marcel Dekker, New York, New York. 314 pp.
- Meyer, D.L., M.S. Fonseca, P.L. Murphey, R.H. McMichael, Jr., M.M. Byerly, M.W. LaCroix, P.E. Whitfield, and G.W. Thayer. 1999. Effects of live-bait shrimp trawling on seagrass beds and fish by-catch in Tampa Bay, Florida. *Fishery Bulletin* 97:193–199.
- Meyer, H.A. and S.S. Bell. 1989. Response of harpacticoid copepods to detrital accumulation on seagrass blades: a field experiment with *Metis holothuriae* (Edwards). *Journal of Experimental Marine Biology and Ecology* 132:141–149.
- Miller, R.L. and B.F. McPherson. 1995. Modeling photosynthetically active radiation in waters of Tampa Bay, Florida, with emphasis on the geometry of incident irradiance. *Estuarine, Coastal and Shelf Science* 40:359–377.
- Milon, J. and E.M. Thunberg. 1993. A regional analysis of current and future Florida resident participation in marine recreational fishing. SGR-112. Florida Sea Grant College Program, Gainesville, Florida.
- Moffler, M.D. and M.J. Durako. 1987. Reproductive biology of the tropical-subtropical seagrasses of the southeastern United States. Pp. 77–88 in Durako, M.J., R.C. Phillips, and R.R. Lewis III (eds.), Proceedings of the Symposium on Subtropical-Tropical Seagrasses of the Southeastern United States. Florida Marine Research Publications No. 42. Florida Marine

- Research Institute, St. Petersburg, Florida. 209 pp. Moncreiff, C.A. and M.J. Sullivan. 2001. Trophic importance of epiphytic algae in subtropical seagrass beds: evidence from multiple stable isotope analyses. *Marine Ecology Progress Series* 215:93–106.
- Montague, C.L. and J.A. Ley. 1993. A possible effect of salinity fluctuation on abundance of benthic vegetation and associated fauna in northeastern Florida Bay. *Estuaries* 16:703–717.
- Montague, C.L. and R.G. Wiegert. 1990. Salt marshes. Pp. 481–516 *in* Myers, R.L. and J. J. Ewel (eds.), *Ecosystems of Florida*. University of Central Florida Press, Orlando, Florida. 765 pp.
- Montague, C.L., A.V. Zale, and H.F. Percival. 1987. The nature of export from fringing marshes with reference to the production of estuarine animals and the effect of impoundments. Pp. 437–450 in Whitman, W.R. and W.H. Meredith (eds.), Waterfowl and Wetlands Symposium: Proceedings of the Symposium on Waterfowl and Wetlands Management in the Coastal Zone of the Atlantic Flyway. Delaware Coastal Management Program, Department of Natural Resources and Environmental Control, Dover, Delaware.
- Moriarty, D.J.W. and P.I. Boon. 1989. Interactions of seagrasses with sediment and water. Pp. 500–535 in Larkum, A.W.D., A.J. McComb, and S. A. Shepherd (eds.), Biology of Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region. Elsevier, Amsterdam, The Netherlands. 841 pp.
- Morris, L.J. and D.A. Tomasko (eds.). 1993. Proceedings and Conclusions of Workshops on Submerged Aquatic Vegetation Initiative and Photosynthetically Active Radiation. Special Publication SJ93-SP13. St. Johns River Water Management District, Palatka, Florida. 244 pp.
- Morrison, G., N. Holland, and H. Greening. 2003a. Conserving Florida's seagrass resources: developing a coordinated statewide management program. Report to the Florida Department of Environmental Protection, Florida Coastal Management Program pursuant to National Oceanic and Atmospheric Administration Award No. NA17OZ2330. FMRI File Code F2361-02. Florida Fish and Wildlife Conservation Commission, Florida Marine Research Institute, St. Petersburg, Florida. 39 pp.
- Morrison, G., A. Janicki, D. Wade, J. Martin, G. Vargo, and R. Johansson. 1997. Estimated nitrogen fluxes and nitrogen-chlorophyll relationships in Tampa

- Bay, 1985–1994. Pp. 249–268 in Treat, S.F. (ed.), Proceedings, Tampa Bay Area Scientific Information Symposium 3: Applying Our Knowledge. Tampa Bay Estuary Program, St. Petersburg, Florida. 396 pp.
- Morrison, G., W. Sargent, and K. Madley. 2003b. Florida seagrass manager's toolkit. Report to the Florida Department of Environmental Protection, Florida Coastal Management Program pursuant to National Oceanic and Atmospheric Administration Award No. NA17OZ2330. FMRI File Code F2361-02. Florida Fish and Wildlife Conservation Commission, Florida Marine Research Institute, St. Petersburg, Florida.
- Morton, R.D., T.W. Duke, J.M. Macauley, J.R. Clark, W.A. Price, S.J. Hendricks, S.L. Owsley-Montgomery, and G.R. Plaia. 1986. Impact of drilling fluids on seagrasses: an experimental community approach. Pp. 199–212 in Cairns, J., Jr. (ed.), Community Toxicity Testing. ASTM STP 920. American Society for Testing and Materials, Philadelphia, Pennsylvania. 350 pp.
- Murali, R.S. 1982. Zero-energy coast. Pp. 883 in Swartz, M.L. (ed.), *The Encyclopedia of Beaches and Coastal Environments*. Hutchinson and Ross Publishers, Stroudsburg, Pennsylvania. 940 pp.
- National Marine Fisheries Service (NMFS). 2002. Recovery plan for Johnson's Seagrass (*Halophila johnsonii*). National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Washington, D.C.
- Neely, M.B. 2000. Somatic, respiratory, and photosynthetic responses of the seagrass *Halodule wrightii* to light reduction in Tampa Bay, Florida including a whole plant carbon budget. Pp. 34–48 *in* Bortone, S.A (ed.), *Seagrasses: Monitoring, Ecology, Physiology, and Management.* CRC Press, Boca Raton, Florida. 318 pp.
- Nelson, W.G. 1980. A comparative study of amphipods in seagrasses from Florida to Nova Scotia. *Bulletin of Marine Science* 30:80–89.
- Nielsen, S.L., I. Thingstrup, and C. Wigand. 1999. Apparent lack of vesicular-arbuscular mycorrhiza (VAM) in the seagrasses *Zostera marina* L. and *Thalassia testudinum* Banks ex König. *Aquatic Botany* 63:261–266.
- Norkko, J., E. Bonsdorff, and A. Norkko. 2000. Drifting algal mats as an alternative habitat for benthic invertebrates: species-specific responses to a transient resource. *Journal of Experimental Marine Biology and Ecology* 248:79–104.
- Norton, T.A. and A.C. Mathieson. 1983. The biology

- of unattached seaweeds. *Progress in Phycological Research* 2:333–386.
- Odum, W.E. and C.C. McIvor. 1990. Mangroves. Pp. 517–548 *in* Myers, R.L. and J. J. Ewel (eds.), *Ecosystems of Florida*. University of Central Florida Press, Orlando, Florida. 765 pp.
- Ogden, J.C. 1980. Faunal relationships in Caribbean seagrass beds. Pp. 173–198 *in* Phillips, R.C. and C.P. McRoy (eds.), *Handbook of Seagrass Biology—An Ecosystem Perspective*. Garland STPM Press, New York, New York. 353 pp.
- Ogden, J.C. 1992. The impact of Hurricane Andrew on the ecosystems of south Florida. *Conservation Biology* 6:488–490.
- O'Leary, M.H. 1988. Carbon isotopes in photosynthesis. *BioScience* 38:328–336.
- Onuf, C.P. 1994. Seagrasses, dredging and light in Laguna Madre, Texas, U.S.A. Estuarine, Coastal and Shelf Science 39:75–91.
- Opsahl, S. and R. Benner. 1993. Decomposition of senescent blades of the seagrass *Halodule wrightii* in a subtropical lagoon. *Marine Ecology Progress Series* 94:191–205.
- Orth, R.J., K.L. Heck, and J. van Montfrans. 1984. Faunal communities in seagrass beds: a review of the influence of plant structure and prey characteristics on predator-prey relationships. *Estuaries* 7:339–350.
- Orth, R.J. and K.A. Moore. 1983. Chesapeake Bay: an unprecedented decline in submerged aquatic vegetation. *Science* 222:51–53.
- Patriquin, D.G. and R. Knowles. 1972. Nitrogen fixation in the rhizosphere of marine angiosperms. *Marine Biology* 16:49–58.
- Phillips, R.C. 1960a. Observations on the ecology and distribution of the Florida seagrasses. *Marine Research Laboratory Professional Papers Series No. 2.* Florida State Board of Conservation, St. Petersburg, Florida. 72 pp.
- Phillips, R.C. 1960b. Ecology and distribution of marine algae found in Tampa Bay, Boca Ciega Bay, and at Tarpon Springs, Florida. *Quarterly Journal of the Florida Academy of Science* 23:23–260.
- Phillips, R.C. 1960c. The ecology of marine plants of Crystal Bay, Florida. Quarterly Journal of the Florida Academy of Science 23:328–337.
- Phillips, R.C. 1984. The ecology of eelgrass meadows in the Pacific Northwest: a community profile. USFWS/OBS-84/24. U.S. Fish and Wildlife Service, Washington, D.C. 85 pp.

- Phlips, E.J., J. Ihnat, and M. Conroy. 1992. Nitrogen fixation by the benthic freshwater cyanobacterium, *Lyngbya wollei*. *Hydrobiologia* 234:59–64.
- Porter, D. and L.K. Muehlstein. 1989. A species of Labyrinthula is the prime suspect as the cause of a massive die-off of the seagrass Thalassia testudinum in Florida Bay. Mycological Society of America Newsletter 40:43.
- Powell, G.V.N., W.J. Kenworthy, and J. W. Fourqurean. 1989. Experimental evidence for nutrient limitation of seagrass growth in a tropical estuary with restricted circulation. *Bulletin of Marine Science* 44:324–340.
- Prager, E.J. and R.B. Halley. 1999. The influence of seagrass on shell layers and Florida Bay mudbanks. *Journal of Coastal Research* 15:1151–1162.
- Pribble, J.R., A.J. Janicki, and H. Greening (eds.). 1999. Baywide environmental monitoring report, 1993–1998: Tampa Bay, Florida. *Tampa Bay Estuary Program Technical Publication #07-99*. Tampa Bay Estuary Program, St. Petersburg, Florida.
- Pulich, W. M. 1986. Variations in leaf soluble amino acids and ammonium content in subtropical seagrasses related to salinity stress. *Plant Physiology* 80:283–286.
- Quammen, M.L. and C.P. Onuf. 1993. Laguna Madre: seagrass changes continue decades after salinity reduction. *Estuaries* 16:302–310.
- Rey, J.R. and F.C. Stephens. 1996. Effects of shading and rhizome isolation on soluble carbohydrate levels in blades and rhizomes of the seagrass Syringodium filiforme. Gulf of Mexico Science 2:47–54.
- Robblee, M.B., T.R. Barber, P.R. Carlson, Jr., M.J. Durako, J.W. Fourqurean, L.K. Muehlstein, D. Porter, L.A. Yarbro, R.T. Zieman, and J.C. Zieman. 1991. Mass mortality of the tropical seagrass *Thalassia testudinum* in Florida Bay (USA). *Marine Ecology Progress Series* 71:297–299.
- Rose, C.D. and C.J. Dawes. 1999. Effects of community structure on the seagrass *Thalassia testudinum*. *Marine Ecology Progress Series* 184:83–95.
- Rose, C.D. and M.J. Durako. 1994. Induced photomorphogenesis by an altered R:FR light ratio in axenic *Ruppia maritima* L. *Botanica Marina* 37:531–535.
- Rose, C.D., W.C. Sharp, W.J. Kenworthy, J.H. Hunt, W.G. Lyons, E.J. Prager, J.F. Valentine, M.O. Hall, P.E. Whitfield, and J.W. Fourqurean. 1999. Overgrazing of a large seagrass bed by the sea urchin *Lytechinus*

- variegatus in Outer Florida Bay. Marine Ecology Progress Series 190:211–222.
- Rydene, D.A. and R.E. Matheson, Jr. 2003. Diurnal fish density in relation to seagrass and drift algae cover in Tampa Bay, Florida. *Gulf of Mexico Science* 2003:35–58.
- Saloman, C.H., S.P. Naughton, and J.L. Taylor. 1982. Benthic faunal assemblages of shallow water sand and seagrass habitats, St. Andrew Bay, Florida. U.S. Fish and Wildlife Service, Division of Ecological Services, Panama City, Florida. 22 pages.
- Sand-Jensen, K. 1977. Effect of epiphytes on eelgrass photosynthesis. *Aquatic Botany* 3:55–63.
- Sand-Jensen, K. 1990. Epiphyte shading: its role in resulting depth distribution of submerged aquatic macrophytes. *Folia Geobotanica et Phytotaxonomica* 25:315–320.
- Santos, S.L. and J.L. Simon. 1974. Distribution and abundance of the polychaetous annelids in a south Florida estuary. *Bulletin of Marine Science* 24:669–689.
- Sarasota Bay National Estuary Program (SBNEP). 1995. The voyage to paradise reclaimed: the comprehensive conservation and management plan for Sarasota Bay, Florida. Sarasota Bay National Estuary Program. Sarasota, Florida.
- Sargent, F.J., T.J. Leary, D.W. Crewz, and C.R. Kruer. 1995. Scarring of Florida's seagrasses: assessment and management options. *Florida Marine Research Institute Technical Report TR-1*. Florida Marine Research Institute, St. Petersburg, Florida. 43 pp. plus appendix.
- Schlueter, M.A. and S.I. Guttman. 1998. Gene flow and genetic diversity of turtle grass, *Thalassia testudinum* Banks ex König, in the lower Florida Keys. *Aquatic Botany* 61:147–164.
- Schoellhamer, D.H. 1991. Tributary storm runoff and vessel-generated sediment resuspension at the Port of St. Petersburg, Florida. Pp. 429–437 in Treat, S.F. and P.A. Clark (eds.), Proceedings, Tampa Bay Area Scientific Information Symposium 2: The Watershed. TEXT, Tampa, Florida. 528 pp.
- Setchell, W.A. 1920. Geographical distribution of the marine spermatophytes. *Bulletin of the Torrey Botanical Club* 47:563–579.
- Setchell, W.A. 1929. Morphological and phenological notes on *Zostera marina* L. University of California *Publications in Botany* 14:389–452.
- Setchell, W.A. 1934. Marine plants and Pacific paleogeography. Pp. 3117–3131 *in* Pacific Science Asso-

- ciation and National Research Council of Canada (eds.), *Proceedings of the Fifth Pacific Science Congress*. University of Toronto Press, Toronto, Canada.
- Setchell, W.A. 1935. Geographic elements of the marine flora of the north Pacific Ocean. *American Naturalist* 69:560–577.
- Shafer, D.J. 1999. The effects of dock shading on the seagrass *Halodule wrightii* in Perdido Bay, Alabama. *Estuaries* 22:936–943.
- Sheridan, P.F. 1992. Comparative habitat utilization by estuarine macrofauna within the mangrove ecosystem of Rookery Bay, Florida. *Bulletin of Marine Science* 50:21–39.
- Sheridan, P.F. and R.J. Livingston. 1983. Abundance and seasonality of infauna and epifauna inhabiting a *Halodule wrightii* meadow in Apalachicola Bay, Florida. *Estuaries* 6:407–419.
- Short, F.T., D.M. Burdick, S. Granger, and S.W. Nixon. 1996. Long-term decline in eelgrass, *Zostera marina* L., linked to increased housing development. Pp. 291–298 *in* Kuo, J., R.C. Phillips, D.I. Walker, and H. Kirkman (eds.), *Seagrass Biology: Proceedings of an International Workshop, Rottnest Island* (2 vols.). University of Western Australia, Nedlands, Western Australia. 663 pp.
- Short, F.T., D.M. Burdick, C.A. Short, R.C. Davis, and P.A. Morgan. 2000. Developing success criteria for restored eelgrass, salt marsh and mud flat habitats. *Ecological Engineering* 15:239–252.
- Short, F.T., W.C. Dennison, and D.G. Capone. 1990. Phosphorus-limited growth of the tropical seagrass Syringodium filiforme in carbonate sediments. Marine Ecology Progress Series 62:169–174.
- Short, F.T. and C.M. Duarte. 2001. Methods for the measurement of seagrass growth and production. Pp. 155–182 *in* Short, F.T. and R.G. Coles (eds.), *Global Seagrass Research Methods*. Elsevier Science B.V., Amsterdam, The Netherlands. 473 pp.
- Short, F.T., J. Montgomery, C.F. Zimmermann, and C.A. Short. 1993. Production and nutrient dynamics of a *Syringodium filiforme* Kütz. seagrass bed in Indian River Lagoon, Florida. *Estuaries* 16:323–334.
- Short, F.T. and H.A. Neckles. 1999. The effects of global climate change on seagrasses. *Aquatic Botany* 63:169–196.
- Short, F.T. and S. Wyllie-Echeverria. 1996. Natural and human—induced disturbance of seagrasses. *Environmental Conservation* 23:17–27.
- Smith, K. 1998. A summary of limited vessel entry zones for managed seagrass protection in Florida.

- Florida Department of Environmental Protection, Division of Marine Resources, Bureau of Protected Species Management, Tallahassee, Florida. 30 pp.
- Smith, K. and R. Mezich. 1999. Comprehensive assessment of the effects of single family docks on seagrass in Palm Beach County, Florida. Draft Report for the Florida Fish and Wildlife Commission, Tallahassee, Florida. 22 pp.
- Sogard, S.M., G.V.N. Powell, and J.G. Holmquist. 1987. Epibenthic fish communities on Florida Bay banks: relations with physical parameters and seagrass cover. *Marine Ecology Progress Series* 40:25–39.
- Staugler, E. and J. Ott. 2000. Establishing baseline seagrass health using fixed transects in Charlotte Harbor, Florida. Florida Department of Environmental Protection, Charlotte Harbor Aquatic Preserves, Pine Island, Florida. 58 pp.
- Stedman, S.-M. and J. Hanson. 1997. Habitat connections: wetlands, fisheries, and economics in the Gulf of Mexico coastal states. National Oceanic and Atmospheric Administration, Office of Habitat Conservation, Silver Spring, Maryland.
- Stoner, A.W. 1980. Abundance, reproductive seasonality and habitat preferences of amphipod crustaceans in seagrass meadows of Apalachee Bay, Florida. *Contributions in Marine Science* 23:63–77.
- Stoner, A.W. 1983. Distribution of fishes in seagrass meadows: role of macrophyte biomass and species composition. *Fishery Bulletin* 81:837–846.
- Stoner, A.W., H.S. Greening, J.D Ryan, and R.J. Livingston. 1983. Comparison of macrobenthos collected with cores and suction sampler in vegetated and unvegetated marine habitats. *Estuaries* 6:76–82.
- Stowers, J.F., E. Fehrmann, and A. Squires. 2002. Seagrass scarring in Tampa Bay: impact analysis and management options. Pp. 47–54 in Greening, H.S. (ed.), Proceedings, Seagrass Management: It's Not Just Nutrients! Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Tampa Bay National Estuary Program (TBNEP). 1996. Charting the course for Tampa Bay. Tampa Bay Estuary Program (TBEP) and the U.S. Environmental Protection Agency, Region IV, St. Petersburg, Florida and Atlanta, Georgia.
- Tampa Bay Regional Planning Council (TBRPC). 1984. The future of Tampa Bay. Tampa Bay Regional Planning Council, St. Petersburg, Florida. 270 pp.
- Taylor, W.R. 1928. The marine algae of Florida, with special reference to the Dry Tortugas. Papers from the Tortugas Laboratory of the Carnegie Institution

- of Washington 25. 219 pp., 3 Figures, 37 plates.
- Thayer, G.W., D.R. Colby, and W.F. Hettler, Jr. 1987. Utilization of the red mangrove prop root habitat by fishes in south Florida. *Marine Ecology Progress Series* 35:25–38.
- Thayer, G.W., M.S. Fonseca, and W.J. Kenworthy. 1997. Ecological value of seagrasses: a brief summary for the ASMFC Habitat Committee's SAV subcommittee. Pp. 5–10 in Stephan, C. D. and T. E. Bigford (eds.), Atlantic Coastal Submerged Aquatic Vegetation: A Review of its Ecological Role, Anthropogenic Impacts, State Regulation and Value to Atlantic Coastal Fisheries. ASMFC Habitat Management Series, No. 1. Atlantic States Marine Fisheries Commission, Washington, D.C.
- Thayer, G.W., W.J. Kenworthy, and M.S. Fonseca. 1984. The ecology of eelgrass meadows of the Atlantic coast: a community profile. FWS/OBS-84/02. U.S. Fish and Wildlife Service, Washington, D.C. 147 pp.
- Thayer, G.W., P.L. Murphy, and M.W. LaCroix. 1994. Responses of plant communities in western Florida Bay to the die-off of seagrasses. *Bulletin of Marine Science* 54:718–726.
- Thayer, G.W., A.B. Powell, and D.E. Hoss. 1999. Composition of larval, juvenile and small adult fishes relative to changes in environmental conditions in Florida Bay. *Estuaries* 22:518–533.
- Thayer, G.W., H.H. Stuart, W.J. Kenworthy, J.F. Ustach, and A.B. Hall. 1978. Habitat values of salt marshes, mangroves, and seagrasses for aquatic organisms. Pp. 235–247 in Greeson, P.E., J.R. Clark, and J.E. Clark (eds.), Wetland Functions and Values: The State of Our Understanding. Proceedings, National Symposium on Wetlands. American Water Resources Association, Minneapolis, Minnesota.
- Thayer, G.W. and J.F. Ustach. 1981. Gulf of Mexico wetlands: value, state of knowledge, and research needs. Pp. 1–30 in Proceedings of the Symposium on Environmental Research Needs in the Gulf of Mexico (GOMEX) 1979, Volume IIB. Key Biscayne, Florida.
- Thomas, M.H. and N. Stratis. 2001. Assessing the economic impact and value of Florida's public piers and boat ramps. Florida Fish and Wildlife Conservation Commission, Tallahassee, Florida. 22 pp.
- Tomasko, D.A. 1992. Variation in growth form of shoal grass (*Halodule wrightii*) due to changes in the spectral composition of light below a canopy of turtle grass (*Thalassia testudinum*). Estuaries

- 15:214-217.
- Tomasko, D.A. 2002. Status and trends of seagrass coverage in Tampa Bay, with reference to other southwest Florida estuaries. Pp.11–20 in Greening, H.S. (ed.), Proceedings, Seagrass Management: It's Not Just Nutrients! Tampa Bay Estuary Program, St. Petersburg, Florida. 246 pp.
- Tomasko, D.A., M. Alderson, P. Clark, J. Culter, K. Dixon, R. Edwards, E. Estevez, M. Heyl, S. Lowery, Y.P. Sheng, and J. Stevely. 1992. Technical synthesis of Sarasota Bay. Pp. 14.1–14.16 in Roat, P., C. Ciccolella, H. Smith, and D.A. Tomasko (eds.), Sarasota Bay: Framework for Action. Sarasota Bay National Estuary Program, Sarasota, Florida.
- Tomasko, D.A., D.L. Bristol, and J.A. Ott. 2001. Assessment of present and future nitrogen loads, water quality, and seagrass (*Thalassia testudinum*) depth distribution in Lemon Bay, Florida. *Estuaries* 24:926–938.
- Tomasko, D.A. and C.J. Dawes. 1989. Evidence for physiological integration between shaded and unshaded short shoots of *Thalassia testudinum*. *Marine Ecology Progress Series* 54:299–305.
- Tomasko, D.A., C.J. Dawes, and M.O. Hall. 1996. The effects of anthropogenic nutrient enrichment on turtle grass, *Thalassia testudinum*, in Sarasota Bay, Florida. *Estuaries* 19:448–456.
- Tomasko, D.A. and K.H. Dunton. 1991. Growth and production of *Halodule wrightii* in relation to continuous measurements of underwater light levels in south Texas. Pp. 79–84 in Kenworthy W.J. and D.E. Haunert (eds.), *The Light Requirements of Seagrasses: Proceedings of a Workshop to Examine the Capability of Water Quality Criteria, Standards, and Monitoring Programs to Protect Seagrasses.* National Oceanic and Atmospheric Administration Technical Memorandum, NMFS-SEFC-287. National Oceanic and Atmospheric Administration, Washington, D.C.
- Tomasko, D.A. and K.H. Dunton. 1995. Primary productivity in *Halodule wrightii*: a comparison of techniques based on daily carbon budgets. *Estuaries* 18:271–278.
- Tomasko, D.A. and M.O. Hall. 1999. Productivity and biomass of the seagrass *Thalassia testudinum* along a gradient of freshwater influence in Charlotte Harbor, Florida. *Estuaries* 22:592–602.
- Tomasko, D.A. and B.E. Lapointe. 1991. Productivity and biomass of *Thalassia testudinum* as related to water column nutrient availability and epiphyte

- levels: field observations and experimental studies. *Marine Ecology Progress Series* 75:9–17.
- Tomlinson, P.B. 1974. Vegetative morphology and meristem dependence—The foundation of productivity in seagrasses. *Aquaculture* 4:107–130.
- Tomlinson, P.B. 1980. Leaf morphology and anatomy in seagrasses. Pp. 7–28 *in* Phillips, R.C. and C.P. McRoy (eds.), *Handbook of Seagrass Biology—An Ecosystem Perspective*. Garland STPM Press, New York. 353 pp.
- Tomlinson, P.B. 1982. Anatomy of the Monocotyledons. Volume VII. Helobiae (Alismatidae). Metcalfe, C.R. (series ed.), Clarendon Press, Oxford, England. 559 pp.
- Touchette, B.W. and J.M. Burkholder. 2000. Review of nitrogen and phosphorus metabolism in seagrasses. *Journal of Experimental Marine Biology and Ecology* 250:133–167.
- Treat, S.F. (ed.). 1997. Proceedings, Tampa Bay Area Scientific Information Symposium 3: Applying our Knowledge. Tampa Bay Regional Planning Council, Tampa, Florida. 396 pp.
- Treat, S.F. and P.A. Clark (eds.). 1991. Proceedings, Tampa Bay Area Scientific Information Symposium 2: The Watershed. TEXT, Tampa, Florida. 528 pp.
- Trocine, R.P., J.D. Rice, and G.N. Wells. 1982. Photosynthetic response of seagrasses to ultraviolet-A radiation and the influence of visible light intensity. *Plant Physiology* 69:341–344.
- Uhrin, A.V. and J.G. Holmquist. 2003. Effects of propeller scarring on macrofaunal use of the seagrass *Thalassia testudinum. Marine Ecology Progress Series* 250:61–70.
- Valentine, J.F. and K.L Heck, Jr. 1993. Mussels in seagrass meadows: their influence on macroinvertebrate abundance and secondary production in the northern Gulf of Mexico. *Marine Ecology Progress Series* 96:63–74.
- Valentine, J.F. and K.L. Heck, Jr. 1999. Seagrass herbivory: evidence for the continued grazing of marine grasses. *Marine Ecology Progress Series* 176:291–302.
- Valentine, J.F., K.L. Heck, Jr., J. Busby, and D. Webb. 1997. Experimental evidence that herbivory increases shoot density and productivity in a subtropical turtlegrass (*Thalassia testudinum*) meadow. *Oecologia* 112:193–200.
- Valentine, J.F., K.L. Heck, Jr., K.D. Kirsch and D. Webb. 2000. Role of sea urchin *Lytechinus variegatus* grazing in regulating subtropical turtlegrass *Thalassia testudinum* meadows in the Florida Keys

- (USA). Marine Ecology Progress Series 200:213–228. van Tussenbroek, B.I. 1994. Aspects of reproductive ecology of *Thalassia testudinum* in Puerto Morelos Reef Lagoon, Mexico. *Botanica Marina* 37:413–419.
- van Tussenbroek, B.I. 1996a. Integrated growth patterns of turtle grass, *Thalassia testudinum* Banks ex König. *Aquatic Botany* 55:139–144.
- van Tussenbroek, B.I. 1996b. Techniques of rapid assessment of seagrass production yield, applied to *Thalassia testudinum* in a Mexican tropical reef lagoon. Pp.131–138 *in* Kuo, J., R.C. Phillips, D.I. Walker, and H. Kirkman (eds.), *Seagrass Biology: Proceedings of an International Workshop, Rottnest Island* (2 vols.). University of Western Australia, Nedlands, Western Australia. 663 pp.
- Virnstein, R.W. 1982. Leaf growth rate of the seagrass *Halodule wrightii* photographically measured *in situ*. *Aquatic Botany* 12:209–218.
- Virnstein, R.W. 1987. Seagrass-associated invertebrate communities of the southeastern USA: a review. Pp. 89–116 in Durako, M.J., R.C. Phillips, and R.R. Lewis III (eds.), Proceedings of the Symposium on Subtropical-Tropical Seagrasses of the Southeastern United States. Florida Marine Research Publications No. 42. Florida Marine Research Institute, St. Petersburg, Florida. 209 pp.
- Virnstein, R.W. 1999. Seagrass meadows: fish and wildlife factories. *Florida Naturalist* 72:18–19.
- Virnstein, R.W. and P.K. Howard. 1987. Motile epifauna of marine macrophytes in the Indian River Lagoon, Florida. II. Comparisons between drift algae and three species of seagrasses. *Bulletin of Marine Science* 41:13–26.
- Virnstein, R.W. and L.J. Morris. 1996. Seagrass preservation and restoration: a diagnostic plan for the Indian River Lagoon. *Technical Memorandum 14*. St. Johns River Water Management District, Palatka, Florida. 43 pp.
- Walters, K. 1991. Influences of abundance, behavior, species composition, and ontogenetic stage on active emergence of meiobenthic copepods in subtropical habitats. *Marine Biology* 108:207–215.
- Walters, K. and S.S. Bell. 1986. Diel patterns of active vertical migration in seagrass meiofauna. *Marine Ecology Progress Series* 34:95–103.
- Waycott, M. and P.A.G. Barnes. 2001. AFLP diversity within and between populations of the Caribbean seagrass *Thalassia testudinum* (Hydrocharitaceae). *Marine Biology* 139:1021–1028.
- Waycott, M., D.W. Freshwater, R.A. York, A. Calladine,

- and W.J. Kenworthy. 2002. Evolutionary trends in the seagrass genus *Halophila* (Thouars): insights from molecular phylogeny. Bulletin of Marine Science 7:1299–1308.
- Wear, D.J., M.J. Sullivan, A.D. Moore, and D.F. Millie. 1999. Effects of water-column enrichment on the production dynamics of three seagrass species and their epiphytic algae. *Marine Ecology Progress Series* 179:201–213.
- Webster, J. and B. A. Stone. 1994. Isolation, structure and monosaccharide composition of the walls of vegetative parts of *Heterozostera tasmanica* (Martens ex Aschers.) den Hartog. *Aquatic Botany* 47:39–52.
- White, C. and J.W. Snodgrass. 1990. Recent changes in the distribution of *Caulerpa prolifera* in the Indian River Lagoon, Florida. *Florida Scientist* 53:85–88.
- Wijesinghe, D. K. and S. N. Handel. 1994. Advantages of clonal growth in heterogeneous habitats: an experiment with *Potentilla simplex*. *Journal of Ecology* 82:495–502.
- Williams, S.L. 1987. Competition between the seagrass Thalassia testudinum and Syringodium filiforme in a Caribbean lagoon. Marine Ecology Progress Series 35:91–98.
- Williams, S.L. 1988a. *Thalassia testudinum* productivity and grazing by green turtles in a highly disturbed seagrass bed. *Marine Biology* 98:447–455.
- Williams, S.L. 1988b. Disturbance and recovery of a deep-water Caribbean seagrass bed. *Marine Ecology Progress Series* 42:63–71.
- Williams, S. L. 1990. Experimental studies of Caribbean seagrass bed development. *Ecological Monographs* 60:449–469.
- Wingrove, P. 1999. Seagrass outreach tool box. University of Florida/Monroe County Cooperative Extension. Florida Bay Education Project, Key Largo, Florida.
- Witz, M.J.A. 1994. Flowering frequency, sex distribution, density and age of short shoots in three west-central Florida *Thalassia testudinum* Banks ex König (Hydrocharitaceae) meadows. M.S. Thesis. University of South Florida, Tampa, Florida.
- Witz, M. J. A. and C. J. Dawes. 1995. Flowering and short shoot age in three *Thalassia testudinum* meadows off west-central Florida. *Botanica Marina* 38:431–436.
- Zarbock, H.W. 1991. Past, present and future freshwater inflow to Tampa Bay: effects of a changing watershed. Pp. 23–34 *in* Treat, S.F. and P.A. Clark

- (eds.), Proceedings, Tampa Bay Area Scientific Information Symposium 2: The Watershed. TEXT, Tampa, Florida. 528 pp.
- Zarbock, H.W., A.J. Janicki, D.L. Wade, D. Haimbuch, and H. Wilson. 1994. Estimates of total nitrogen, total phosphorus, and total suspended solids loadings to Tampa Bay, Florida. *Tampa Bay Estuary Program Technical Publication #04-94*. Tampa Bay Estuary Program, St. Petersburg, Florida.
- Zarbock, H.W., A.J. Janicki, D.L. Wade, and R.J. Pribble. 1996. Model-based estimates of total nitrogen loading to Tampa Bay. *Tampa Bay Estuary Program Technical Publication #05-96*. Tampa Bay Estuary Program, St. Petersburg, Florida.
- Zarbock, H.W., J. Schulten, E. Long, and D. Mac-Donald. 1997. Sediment contamination in Tampa Bay: sources, risks and management. Pp. 281–293 in Treat, S.F. (ed.), Proceedings, Tampa Bay Area Scientific Information Symposium 3: Applying Our Knowledge. Tampa Bay Estuary Program, St. Petersburg, Florida. 396 pp.
- Zieman, J.C. 1976. The ecological effects of physical damage from motor boats on turtle grass beds in southern Florida. *Aquatic Botany* 2:127–139.
- Zieman, J.C. 1982. The ecology of the seagrasses of south Florida: a community profile. FWS/OBS-82/25. U.S. Fish and Wildlife Services, Washington, D.C. 158 pp.
- Zieman, J.C. 1987. A review of certain aspects of the life, death, and distribution of the seagrasses of the southeastern United States, 1960–1985. Pp. 53–76 in Durako, M.J., R.C. Phillips, and R.R. Lewis III (eds.), Proceedings of the Symposium on Subtropical-Tropical Seagrasses of the Southeastern United States. Florida Marine Research Publications No. 42. Florida Marine Research Institute, St. Petersburg, Florida. 209 pp.
- Zieman, J.C., J.W. Fourqurean, and T.A. Frankovich. 1999. Seagrass die-off in Florida Bay (U.S.A.): longterm trends in abundance and growth of turtle grass, *Thalassia testudinum*. *Estuaries* 22:460–470.
- Zieman, J.C., J.W. Fourqurean, and R. L. Iverson. 1989. Distribution, abundance and productivity of seagrasses and macroalgae in Florida Bay. *Bulletin of Marine Science* 44: 292–311.
- Zieman, J.C. and R.G. Wetzel. 1980. Productivity in seagrasses: methods and rates. Pp. 87–116 in Phillips, R.C. and C.P. McRoy (eds.), Handbook of Seagrass Biology—An Ecosystem Perspective. Garland STPM Press, New York, New York. 353 pp.

Zieman, J.C. and R. T. Zieman. 1989. The ecology of the seagrass meadows of the west coast of Florida: a community profile. Biological Report 85(7.25). U.S. Fish and Wildlife Service, Washington, D.C. 155 pp. Zimmerman, M.S. and R.J. Livingston. 1976. Effects of kraft-mill effluent on benthic macrophyte assemblages in a shallow-bay system (Apalachee Bay, North Florida, USA). *Marine Biology* 34: 297–312. Zimmerman, R.C., A. Cabello-Pasini, and R.S. Alberte. 1994. Modeling daily production of aquatic macrophytes from irradiance measurement: a comparative analysis. *Marine Ecology Progress Series* 114:185–196.

